

REFERENCES

- Adin, R. & Riera, P. (2003) Preferential food source utilization among stranded macroalgae by *Talitrus saltator* (Amphipoda, Talitridae): a stable isotopes study in the northern coast of Brittany (France), *Estuarine Coastal and Shelf Science* **56**, 91-98.
- Alongi, D. M. (1998) Beaches and tidal flats, *In: Coastal ecosystem processes*, CRC Press, New York.
- Barros, F., Underwood, A. J. & Archambault, P. (2004) The influence of troughs and crests of ripple marks on the structure of subtidal benthic assemblages around rocky reefs, *Estuarine Coastal and Shelf Science* **60**, 781-790.
- Beaver, B. V., Reed, W., Leary, S., McKiernan, B., Bain, F., Schultz, R., Bennett, B. T., Pascoe, P., Shull, E., Cork, L. C., Francis-Floyd, R., Amass, K. D., Johnson, R., Schmidt, R. H., Underwood, W., Thornton, G. W. & Kohn, B. (2000) Report of the AVMA Panel on Euthanasia, *Journal of the American Veterinary Medical Association* **218**, 669-696.
- Behbehani, M. I. & Croker, R. A. (1982) Ecology of beach wrack in northern New England with special reference to *Orchestia platensis*, *Estuarine Coastal and Shelf Science* **15**, 611-620.
- Berg, B. & Laskowski, R. (2006) *Litter Decomposition: A Guide to Carbon and Nutrient Turnover*, Elsevier, Burlington, USA.
- Blanche, K. R. (1992) Preliminary observations on the distribution and abundance of seaweed flies (Diptera: Coleopidae) on beaches in the Gosford district of New South Wales, Australia, *Australian Journal of Ecology* **17**, 27-34.
- Blott, S. J. & Pye, K. (2001) Gradistat: a grain size distribution and statistics package for the analysis of unconsolidated sediments, *Earth Surface Processes and Landforms* **26**, 1237-1248.
- Bosley, K. L. & Wainwright, S. C. (1999) Effects of preservatives and acidification on the stable isotope ratios ($^{15}\text{N}:$ ^{14}N , $^{13}\text{C}:$ ^{12}C) of two species of marine animals, *Canadian Journal of Fisheries and Aquatic Sciences* **56**, 2181-2185.

- Boulton, A. J. & Boon, P. I. (1991) A review of methodology used to measure leaf litter decomposition in lotic environments: time to turn over an old leaf?, *Australian Journal of Marine and Freshwater Research* **42**, 1-43.
- Bradley, R. A. & Bradley, D. W. (1993) Wintering shorebirds increase after kelp (*Macrocystis*) recovery, *The Condor* **95**, 372-376.
- Brown, A. C. & McLachlan, A. (2002) Sandy shore ecosystems and the threats facing them: some predictions for the year 2025, *Environmental Conservation* **29**, 62-77.
- Bryars, S., Miller, D., Collings, G., Fernandes, M., Mount, G. & Wear, R. (2006) Field surveys 2003-2005: Assessment of the quality of Adelaide's coastal waters, sediments and seagrasses. ACWS Technical Report No. 14 prepared for the Adelaide Coastal Waters Study Steering Committee, South Australian Research and Development Institute (Aquatic Sciences), Publication No. RD01/0208-15, Adelaide.
- Bustamante, R. H. & Branch, G. M. (1996) The dependence of intertidal consumers on kelp-derived organic matter on the west coast of South Africa, *Journal of Experimental Marine Biology and Ecology* **196**, 1-28.
- Chapman, M. G. & Roberts, D. E. (2004) Use of seagrass wrack in restoring disturbed Australian saltmarshes, *Ecological management and restoration* **5**, 183-190.
- Clarke, K. R. & Warwick, R. M. (1994) Change in marine communities: An approach to statistical analysis and interpretation, Bourne Press Limited, Bournemouth, UK.
- Colombini, I., Aloia, A., Fallaci, M., Pezzoli, G. & Chelazzi, L. (2000) Temporal and spatial use of stranded wrack by the macrofauna of a tropical sandy beach, *Marine Biology* **136**, 531-541.
- Colombini, I. & Chelazzi, L. (2003) Influence of marine allochthonous input on sandy beach communities, *Oceanography and Marine Biology* **41**, 115-159.
- Connolly, R. M., Hindell, J. S. & Gorman, D. (2005) Seagrass and epiphytic algae support nutrition of a fisheries species, *Sillago schomburgkii*, in adjacent intertidal habitats, *Marine Ecology Progress Series* **286**, 69-79.

- Coupland, G. T. & McDonald, J. I. (2008) Extraordinarily high earthworm abundance in deposits of marine macrodetritus along two semi-arid beaches, *Marine Ecology Progress Series* **361**, 181-9.
- Crawley, K. R., Hyndes, G. A. & Ayvazian, S. G. (2006) Influence of different volumes and types of detached macrophytes on fish community structure in surf zones of sandy beaches, *Marine Ecology Progress Series* **307**, 233-246.
- Cubit, J. (1969) Behaviour and physical factors causing migration and aggregation of the sand crab *Emerita analoga* (Stimpson), *Ecology* **50**, 118-123.
- Currin, C. A., Newell, S. Y. & Paerl, H. W. (1995) The role of standing dead *Spartina alterniflora* and benthic microalgae in salt marsh food webs: considerations based on multiple stable isotope analysis, *Marine Ecology Progress Series* **121**, 99-116.
- Davenport, S. R. & Bax, N. J. (2002) A trophic study of a marine ecosystem off southeastern Australia using stable isotopes of carbon and nitrogen, *Canadian Journal of Fisheries and Aquatic Sciences* **59**, 514-530.
- de Falco, G., Simeone, S. & Baroli, M. (2008) Management of beach-cast *Posidonia oceanica* seagrass on the island of Sardinia (Italy, western Mediterranean), *Journal of Coastal Research* **24**, 69-75.
- Dugan, J. E., Hubbard, D. M., McCrary, M. D. & Pierson, M. O. (2003) The response of macrofauna communities and shorebirds to macrophyte wrack subsidies on exposed sandy beaches of southern California, *Estuarine Coastal and Shelf Science* **58**, 25-40.
- Duggins, D. O. & Eckman, J. E. (1997) Is kelp detritus a good food for suspension feeders? Effects of kelp species, age and secondary metabolites, *Marine Biology* **128**, 489-495.
- Duggins, D. O., Simenstad, C. A. & Estes, J. A. (1989) Magnification of secondary production by kelp detritus in coastal marine ecosystems, *Science* **245**, 170-173.
- Edgar, G. J. (1987) Dispersal of faunal and floral propagules associated with drifting *Macrocystis pyrifera* plants, *Marine Biology* **95**, 599-610.
- Egglisshaw, H. J. (1965) Observations on the fauna of wrack beds, *Transactions of the Society for British Entomology* **16**, 189-216.

- Eleftheriou, A. & McIntyre, A. (2005) *Methods for the Study of Marine Benthos*, Blackwell Science Ltd, Oxford, UK.
- Engelhard, T. & Withers, K. (1998) Biological effects of mechanical beach raking in the upper intertidal zone on Padre Island national seashore, Texas, *GERS Meeting Abstracts 1998*.
- Fairweather, P. G. (1991) Statistical power and design requirements for environmental monitoring, *Australian Journal of Marine and Freshwater Research* **42**, 555-567.
- Fairweather, P. G. & Henry, R. J. (2003) To clean or not to clean? Ecologically sensitive management of wrack deposits on sandy beaches, *Ecological Management and Restoration* **4**, 8-9.
- Fanini, L., Cantarino, C. M. & Scapini, F. (2005) Relationships between the dynamics of two populations of *Talitrus saltator* and the impacts of activities linked to tourism, *Oceanologia* **47**, 93-112.
- Gheskiere, T., Magda, V., Greet, P. & Steven, D. (2006) Are strandline meiofaunal assemblages affected by a once-only mechanical beach cleaning? Experimental findings, *Marine Environmental Research* **61**, 245-264.
- Gimenez, L. & Yannicelli, B. (2000) Longshore patterns of distribution of macroinfauna on a Uruguayan sandy beach: An analysis at different spatial scales and of their potential causes, *Marine Ecology Progress Series* **199**, 111-125.
- Gobert, S., Cambridge, M. L., Velimirov, B., Pergent, G., Lepoint, G., Bouquegneau, J., Dauby, P., Pergent-Martini, C. & Walker, D. I. (2006) *Biology of Posidonia*, In: *Seagrasses: Biology, Ecology and Conservation*, Springer, Dordrecht, The Netherlands.
- Griffiths, C. L. & Stenton-Dozey, J. M. E. (1981) The fauna and rate of degradation of stranded kelp, *Estuarine Coastal and Shelf Science* **12**, 645-653.
- Harrison, P. G. (1982) Control of microbial growth and of amphipod grazing by water-soluble compounds from leaves of *Zostera marina*, *Marine Biology* **67**, 225-230.
- Harrison, P. G. & Mann, K. H. (1975) Detritus formation from eelgrass (*Zostera marina* L.): The relative effects of fragmentation, leaching, and decay, *Limnology and Oceanography* **20**, 924-934.

- Harrold, C., Light, K. & Lisin, S. (1998) Organic enrichment of submarine-canyon and continental-shelf benthic communities by macroalgal drift imported from nearshore kelp forests, *Limnology and Oceanography* **43**, 669-678.
- Harrold, C. & Lisin, S. (1989) Radio-tracking rafts of giant kelp: local production and regional transport, *Journal of Experimental Marine Biology and Ecology* **130**, 237-251.
- Hemminga, M. A. & Nieuwenhuize, J. (1990) Seagrass wrack-induced dune formation on a tropical coast (Banc d'Arguin, Mauritania), *Estuarine Coastal and Shelf Science* **31**, 499-502.
- Hobday, A. J. (2000) Abundance and dispersal of drifting kelp *Macrocystis pyrifera* rafts in the Southern California Bight, *Marine Ecology Progress Series* **195**, 101-116.
- Hubbard, D. M. & Dugan, J. E. (2003) Shorebird use of an exposed sandy beach in southern California, *Estuarine Coastal and Shelf Science* **58**, 41-54.
- Hutchins, B. & Swainston, R. (1986) *Sea Fishes of Southern Australia*, Swainston Publishing, Smithfield, NSW, Australia.
- Ince, R., Hyndes, G. A., Lavery, P. S. & Vanderklift, M. A. (2007) Marine macrophytes directly enhance abundances of sandy beach fauna through provision of food and habitat, *Estuarine Coastal and Shelf Science* **74**, 77-86.
- Inglis, G. (1989) The colonisation and degradation of stranded *Macrocystis pyrifera* (L.) C. Ag. by the macrofauna of a New Zealand sandy beach, *Journal of Experimental Marine Biology and Ecology* **125**, 203-217.
- Jackson, E. L., Rowden, A. A., Attrill, M. J., Bossy, S. F. & Jones, M. B. (2002) Comparison of fish and mobile macroinvertebrates associated with seagrass and adjacent sand at St. Catherine Bay, Jersey (English Channell): Emphasis on commercial species, *Bulletin of Marine Science* **71**, 1333-1341.
- Jacob, U., Mintembeck, K., Brey, T., Knust, R. & Beyer, K. (2005) Stable isotope food web studies: a case for standardised sample treatment, *Marine Ecology Progress Series* **287**, 251-253.
- James, R. (1999) Cusps and pipis on a sandy ocean beach in New South Wales, *Australian Journal of Ecology* **24**, 587-592.

- Jaramillo, E., De la Huz, R., Duarte, C. & Contreras, H. (2006) Algal wrack deposits and macroinfaunal arthropods on sandy beaches of the Chilean coast, *Revista Chilena De Historia Natural* **79**, 337-351.
- Jedrzejczak, M. F. (2002a) Spatio-temporal decay 'hot spots' of stranded wrack in a Baltic sandy coastal system. Part I. Comparative study of the pattern: 1 type of wrack vs 3 beach sites, *Oceanologia* **44**, 491-512.
- Jedrzejczak, M. F. (2002b) Stranded *Zostera marina* L. vs wrack fauna community interactions on a Baltic sandy beach (Hel, Poland): a short-term pilot study. Part I. Driftline effects of fragmented detritivory, leaching and decay rates, *Oceanologia* **44**, 273-286.
- Jedrzejczak, M. F. (2002c) Stranded *Zostera marina* L. vs wrack fauna community interactions on a Baltic sandy beach (Hel, Poland): a short-term pilot study. Part II. Driftline effects of succession changes and colonisation of beach fauna, *Oceanologia* **44**, 367-387.
- Jenkins, G. P. & Wheatley, M. J. (1998) The influence of habitat structure on nearshore fish assemblages in a southern Australian embayment: Comparison of shallow seagrass, reef-algal and unvegetated sand habitats, with emphasis on their importance to recruitment, *Journal of Experimental Marine Biology and Ecology* **221**, 147-172.
- Karban, R. & Huntzinger, M. (2006) How to do ecology, Princeton University Press, Princeton, New Jersey.
- Kendrick, G. A., Kirkman, H. & Burton, C. (1995) An investigation into beach cast macro algae and marine angiosperms, CSIRO Division of Fisheries, Australia.
- Kingsford, M. J. & Choat, J. H. (1985) The fauna associated with drift algae captured with a plankton-mesh purse seine net, *Limnology and Oceanography* **30**, 618-630.
- Kirkman, H. & Kendrick, G. A. (1997) Ecological significance and commercial harvesting of drifting and beach-cast macro-algae and seagrasses in Australia: a review, *Journal of Applied Phycology* **9**, 311-326.
- Koop, K. & Griffiths, C. L. (1982) The relative significance of bacteria, meio- and macrofauna on an exposed sandy beach, *Marine Biology* **66**, 295-300.

- Kuiter, R. H. (1996) A Guide to the Sea Fishes of Australia, New Holland Publishers (Australia) Pty. Ltd., Sydney, Australia.
- Lasiak, T. A. (1986) Juveniles, food and the surf zone habitat: implications for teleost nursery areas *South African Journal of Zoology* **21**, 51-56.
- Lastra, M., Page, H. M., Dugan, J. E., Hubbard, D. M. & Rodil, I. F. (2008) Processing of allochthonous macrophyte subsidies by sandy beach consumers: estimates of feeding rates and impacts on food resources, *Marine Biology* **154**, 163-174.
- Lavery, P., Bootle, S. & Vanderklift, M. (1999) Ecological effects of macroalgal harvesting on beaches in the Peel-Harvey estuary, Western Australia, *Estuarine Coastal and Shelf Science* **49**, 295-309.
- Lavoie, D. R. (1985) Population dynamics and ecology of beach wrack macroinvertebrates of the central California coast, *Bulletin of the Southern Californian Academy of Science* **84**, 1-22.
- Lenanton, R. C. J. & Caputi, N. (1989) The roles of food supply and shelter in the relationship between fishes, in particular *Cnidoglanis macrocephalus* (Valenciennes), and detached macrophytes in the surf zone of sandy beaches, *Journal of Experimental Marine Biology and Ecology* **128**, 165-176.
- Lenanton, R. C. J., Robertson, A. I. & Hansen, J. A. (1982) Nearshore accumulations of detached macrophytes as nursery areas for fish, *Marine Ecology Progress Series* **9**, 51-57.
- Liddle, M. J. & Greig-Smith, P. (1974) A survey of tracks and paths in a sand dune ecosystem. I. Soils, *The Journal of Applied Ecology* **12**, 893-908.
- Llewellyn, P. J. & Shackley, S. E. (1996) The effects of mechanical beach-cleaning on invertebrate populations, *British Wildlife* **70**, 147-155.
- Maccarone, A. D., Brzorad, J. N. & Parsons, K. C. (1993) Nest site selection by herring gulls in an urban estuary, *Colonial Waterbirds* **16**, 216-220.
- Machas, R., Santos, R. & Peterson, B. (2006) Elemental and stable isotope composition of *Zostera noltii* (Horneman) leaves during early phases of decay in a temperate mesotidal lagoon, *Estuarine, Coastal and Shelf Science* **66**, 21-29.

- Malm, T., Raberg, S., Fell, S. & Carlsson, P. (2004) Effects of beach cast cleaning on beach quality, microbial food web, and littoral macrofaunal biodiversity, *Estuarine Coastal and Shelf Science* **60**, 339-347.
- Marsden, I. D. (1991) Kelp sandhopper interactions on a sand beach in New Zealand 1. Drift composition and distribution, *Journal of Experimental Marine Biology and Ecology* **152**, 61-74.
- Masselink, G., Hegge, B. J. & Pattiaratchi, C. B. (1997) Beach cusp morphodynamics, *Earth Surface Processes and Landforms* **22**, 1139-1155.
- Masselink, G. & Hughes, M. G. (2003) Introduction to Coastal Processes & Geomorphology, Hodder Arnold, London.
- Mateo, M. A., Sanchez-Lizaso, J. L. & Romero, J. (2003) *Posidonia oceanica* 'banquettes': a preliminary assessment of the relevance for meadow carbon and nutrients budget, *Estuarine Coastal and Shelf Science* **56**, 85-90.
- McAlpine, D. K. (1991) Review of the Australian kelp flies (Diptera: Coleopidae), *Systematic Entomology* **16**, 29-84.
- McArdle, S. & McLachlan, A. (1991) *Marine Ecology Progress Series* **76**, 91-99.
- McDonald, R. C., Isbell, R. F., Speight, J. G., Walker, J. & Hopkins, M. S. (1990) Australian soil and land survey field handbook, 2nd edition, Inkata Press Pty. Ltd., Melbourne and Sydney, Australia.
- McGwynne, L. E., McLachlan, A. & Furstenberg, J. P. (1988) Wrack breakdown on sandy beaches - Its impact on interstitial meiofauna, *Marine Environmental Research* **25**, 213-232.
- McKechnie, J. M. & Fairweather, P. G. (2003) Ecological Implications for the Management of Wrack on South Australian Sandy Beaches: A Report to the Coast Protection Board, School of Biological Sciences, Flinders University, Adelaide, Australia.
- McLachlan, A. (1981) Exposed sandy beaches as semi-enclosed ecosystems, *Marine Environmental Research* **2**, 59-63.
- McLachlan, A. (1985) The biomass of macro- and interstitial fauna on clean and wrack-covered beaches in Western Australia, *Estuarine Coastal and Shelf Science* **21**, 587-599.

- McLachlan, A. & Brown, A. (2006) The ecology of sandy shores, 2nd edition, Elsevier Inc., Massachusetts, USA.
- McLachlan, A. & Dorvlo, A. (2005) Global patterns in sandy beach macrobenthic communities, *Journal of Coastal Research* **21**, 674-687.
- McLachlan, A. & Dorvlo, A. (2007) Global patterns in sandy beach macrobenthic communities: biological factors, *Journal of Coastal Research* **23**, 1081-1087.
- McLachlan, A. & Hesp, P. (1984) Faunal response to morphology and water circulation of a sandy beach with cusps, *Marine Ecology Progress Series* **19**, 133-144.
- McMillan, C. (1980) $^{13}\text{C}/^{12}\text{C}$ ratios in seagrasses, *Aquatic Botany* **9**, 237-249.
- Mews, M., Zimmer, M. and Jelinski, D.E. (2006) Species-specific decomposition rates of beach-cast wrack in Barkley Sound, British Columbia, Canada, *Marine Ecology Progress Series* **328**, 155-160.
- Nickum, J. G., Bart Jr, H. L., Bowser, P. R., Greer, I. E., Hubbs, C., Jenkins, J. A., MacMillan, J. R., Rachlin, J. W., Rose, J. D., Sorensen, P. W. & Tomasso, J. R. (2004) Guidelines for the Use of Fishes in Research VIII, American Institute of Fishery Research Biologists and American Society of Ichthyologists and Herpetologists.
- Nolan, T. J., Kirk, R. M. & Shulmeister, J. (1999) Beach cusp morphology on sand and mixed sand and gravel beaches, South Island, New Zealand, *Marine Geology* **157**, 185-198.
- Nordstrom, K. F., Lampe, R. & Vandemark, L. M. (2000) Reestablishing naturally functioning dunes on developed coasts, *Environmental Management* **25**, 37-51.
- Nordstrom, K. R., Jackson, N. L., Klein, A. H. F., Sherman, D. J. & Hesp, P. A. (2006) Offshore aeolian transport across a low foredune on a developed barrier island, *Journal of Coastal Research* **22**, 1260-1267.
- Ochieng, C. A. & Erfemeijer, P. L. A. (1999) Accumulation of seagrass beach cast along the Kenyan coast: a quantitative assessment, *Aquatic Botany* **65**, 221-238.
- Olabarria, C., Lastra, M. & Garrido, J. (2007) Succession of macrofauna on macroalgal wrack of an exposed sandy beach: Effects of patch size and site, *Marine Environmental Research* **63**, 19-40.

- Orr, M., Zimmer, M., Jelinski, D. E. & Mews, M. (2005) Wrack deposition on different beach type: spatial and temporal variation in the pattern of subsidy, *Ecology* **86**, 1496-1507.
- Pattiaratchi, C. B., Newgard, J. & Hollings, B. (2007) Physical oceanographic studies of Adelaide coastal waters using high-resolution modelling, in-situ observations and satellite techniques - Sub Task 2 Final Technical Report, ACWS Technical Report No. 20 prepared for the Adelaide Coastal Waters Study Steering Committee, Adelaide.
- Peterson, B. J. & Fry, B. (1987) Stable isotopes in ecosystem studies, *Annual Review of Ecology and Systematics* **18**, 293-320.
- Peterson, C. H., Hickerson, D. H. M. & Grissom Johnson, G. (2000) Short-term consequences of nourishment and bulldozing on the dominant large invertebrates of a sandy beach, *Journal of Coastal Research* **16**, 368-378.
- Phillips, D. S. & Gregg, J. W. (2003) Source partitioning using stable isotopes: coping with too many sources, *Oecologia* **136**, 261-269.
- Piriz, M. L., Eyra, M. C. & Rostagno, C. M. (2003) Changes in biomass and botanical composition of beach-cast seaweeds in a disturbed coastal area from Argentine Patagonia, *Journal of Applied Phycology* **15**, 67-74.
- PIRSA (2003) South Australian Fisheries Management Series. Draft Management Plan for Harvesting Beach-cast Seagrass and Marine Algae, Primary Industries and Resources South Australia, Australia.
- Polis, G. A. & Hurd, S. D. (1996) Linking marine and terrestrial food webs: allochthonous input from the ocean supports high secondary productivity on small islands and coastal land communities, *The American Naturalist* **147**, 396-423.
- Post, D. M. (2002) Using stable isotopes to estimate trophic position: models, methods, and assumptions, *Ecology* **83**, 703-718.
- Raven, J. A., Johnston, A. M., Kubler, J. E., Korb, R., McInroy, S. G., Handley, L. L., Scrimgeour, C. M., Walker, D. I., Beardall, J., Vanderklift, M. A., Fredriksen, S. & Dunton, K. H. (2002) Mechanistic interpretation of carbon isotope

- discrimination by marine macroalgae and seagrasses, *Functional Plant Biology* **29**, 355-378.
- Rieper-Kirchner, M. (1990) Macroalgal decomposition: laboratory studies with particular regard to microorganisms and meiofauna, *Helgolander Meeresuntersuchungen* **44**, 397-410.
- Robertson, A. I. & Hansen, J. A. (1982) Decomposing seaweed: a nuisance of a vital link in coastal food chains?, CSIRO Division of Fisheries Research Report 1980-1981, Australia.
- Robertson, A. I. & Lucas, J. S. (1983) Food choice, feeding rates, and the turnover of macrophyte biomass by a surf-zone inhabiting amphipod, *Journal of Experimental Marine Biology and Ecology* **72**, 99-124.
- Robertson, A. I. & Mann, K. H. (1980) The role of isopods and amphipods in the initial fragmentation of eelgrass detritus in Nova Scotia, Canada, *Marine Biology* **59**, 63-69.
- Rodil, I. F., Olabarria, C., Lastra, M. & Lopez, J. (2008) Differential effects of native and invasive algal wrack on macrofaunal assemblages inhabiting exposed sandy beaches, *Journal of Experimental Marine Biology and Ecology* **358**, 1-13.
- Rodriguez, S. R. (2003) Consumption of drift kelp by intertidal populations of the sea urchin *Tetrapygus niger* on the central Chilean coast: possible consequences at different ecological levels, *Marine Ecology Progress Series* **251**, 141-151.
- Rossi, F. & Underwood, A. J. (2002) Small-scale disturbance and increased nutrients as influences on intertidal macrobenthic assemblages: experimental burial of wrack in different intertidal environments, *Marine Ecology Progress Series* **241**, 29-39.
- Russel, R. J. & McIntire, W. G. (1965) Beach cusps, *Geological Society of America Bulletin* **76**, 307-320.
- Schlacher, T. A., Schoeman, D. S., Dugan, J., Lastra, M., Jones, A., Scapini, F. & McLachlan, A. (2008) Sandy beach ecosystems: key features, sampling issues, management challenges and climate change impacts, *Marine Ecology* **29**, 70-90.
- Short, A. D. (2006a) Australian beach systems - nature and distribution, *Journal of Coastal Research* **22**, 11-27.

- Short, A. D. (2006b) *Beaches of the South Australian Coast and Kangaroo Island*, Sydney University Press, Sydney.
- Short, A. D. & Hesp, P. A. (1982) Wave, beach and dune interactions in south-eastern Australia, *Marine Geology* **48**, 259-284.
- Sousa, W. P. (1979) Disturbance in marine intertidal boulder fields: the nonequilibrium maintenance of species diversity, *Ecology* **60**, 537-546.
- Thayer, G. W., Engel, D. W. & LaCroix, M. W. (1977) Seasonal distribution and changes in the nutritive quality of living, dead and detrital fractions of *Zostera marina* L., *Journal of Experimental Marine Biology and Ecology* **30**, 109-127.
- Thorner, C. S., DiMilla, P., Nixon, S. W. & McKinney, R. A. (2008) Natural and anthropogenic nitrogen uptake by bloom-forming macroalgae, *Marine Pollution Bulletin* **56**, 261-269.
- Underwood, A. J. (1997) *Experiments in Ecology. Their Logical Design and Interpretation Using Analysis of Variance*, Cambridge University Press, Cambridge.
- Urban-Malinga, B., Gheskiere, T., Degraer, S., Derycke, S., Opalinski, K. W. & Moens, T. (2008) Gradients in biodiversity and macroalgal wrack decomposition rate across a macrotidal, ultradissipative sandy beach, *Marine Biology* **155**, 79-90.
- Van der Merwe, D. & McLachlan, A. (1987) Significance of free-floating macrophytes in the ecology of a sandy beach surf-zone, *Marine Ecology Progress Series* **38**, 53-63.
- Vizzini, S. & Mazzola, A. (2006) Sources and transfer of organic matter in food webs of a Mediterranean coastal environment: Evidence for spatial variability, *Estuarine Coastal and Shelf Science* **66**, 459-467.
- Walker, D. I. & McComb, A. J. (1985) Decomposition of leaves from *Amphibolis antarctica* (Labill.) Sonder et Aschers. and *Posidonia australis* Hook. f. the major seagrass species of Shark Bay, Western Australia, *Botanica Marina* **28**, 407-423.
- Wentworth, C. K. (1922) A scale of grade and class terms for clastic sediments, *Journal of Geology* **30**, 377-392.

- Williams, A., Feagin, R. & Witmer, A. (2008) Environmental impacts of beach raking of *Sargassum* spp. on Galveston Island, Texas, *Shore and Beach* **76**, 63-69.
- Womersley, H. B. S. (1984) The Marine Benthic Flora of Southern Australia. Part I, Government Printer, South Australia.
- Womersley, H. B. S. (1987) The Marine Benthic Flora of Southern Australia. Part II, Government Printer, South Australia.
- Womersley, H. B. S. (1994a) The Marine Benthic Flora of Southern Australia. Part IIIA, Australian Biological Resources Study, Canberra.
- Womersley, H. B. S. (1994b) The Marine Benthic Flora of Southern Australia. Part IIIB, Australian Biological Resources Study, Canberra.
- Womersley, H. B. S. (1994c) The Marine Benthic Flora of Southern Australia. Part IIIC, Australian Biological Resources Study, Canberra.
- Yatsuya, K., Nishigaki, T., Douke, A. & Wada, Y. (2007) Seasonal changes in biomass of macrophytes stranded on Yoro Beach along a sargassacean forest in Wakasa Bay, Sea of Japan, *Fisheries Science* **73**, 609-614.
- ZoBell, C. E. (1971) Drift seaweeds on San Diego County beaches, *In: The biology of giant kelp beds (Macrocystis) in California*, North, W.J. (ed), Nova Hedwigia **32**, 269-314.