# **Evolution and Ecology of the Prehensile- Tailed Skink** – *Corucia zebrata*

# Ingerid Julie Hagen MSc

A thesis submitted in fulfilment of the requirements for the degree of Doctor of Philosophy

School of Biological Sciences, Faculty of Engineering, Flinders University

February 2011



## **TABLE OF CONTENTS**

ABSTRACT	5
PUBLICATIONS AND PRESENTATIONS	7
DECLARATION	8
ACKNOWLEDGEMENT	10
CHAPTER 1: GENERAL INTRODUCTION	12
Introduction	13
Island Biogeography and Phylogeography	14
Phylogenetic inference	
Population genetics	17
Field ecology	19
Radio telemetry	
Capture-Mark-Recapture	
Occupance	
Conservation and management	21
Corucia zebrata – life history and ecology as known prior to the current study	22
Corucia in the wider context: the Egernia group and social behaviour	
Properties of the native range: geology and ecotype of the Solomon Archipelago	
Working in the region	
Ugi Island	
Conservation status of C. zebrata	
Project aims and thesis structure	28
CHAPTER 2: PHYLOGEOGRAPHY AND COLONIZATION PATTERN OF CORUCI ZEBRATA ON THE SOLOMON ARCHIPELAGO	30
Introduction	31
Materials and Methods	35
DNA extraction, PCR amplification and sequencing	35
Data analysis	
Results	39
Recombination and statistical properties of the loci	
Phylogenetic relationships between island populations	
Inter-island genetic divergence	
Haplotype networks	
Dating of colonization events within the Solomon Islands	44
Daung of colonization events within the solomon islands	40
Discussion	47
Genetic distance and mitochondrial reciprocal monophyly in relation to Pleistocene land-bridges	
Genetic distance in relation to geographic distance	

Colonization history and dating of colonisation events within the Sc	
Dispersal ability and mechanism of colonisation	
Acknowledgements	52
CHAPTER 3: HANGING ABOUT IN TREES - POPULAT	ON GENETIC STRUCTURE
OF THE PREHENSILE TAILED LIZARD, <i>CORUCIA ZEE</i>	BRATA54
Abstract	55
Introduction	55
Methods and Materials	57
Determination of juvenile vs. adults	
DNA extraction and PCR amplification	
Microsatellite data analysis	
Population sub-structure	62
Individual-level population structure	62
Results	
Captures	
Population sub-structure	
Group structure and relatedness	63
Discussion	66
Acknowledgements	67
_	
CHAPTER 4: HOME RANGES IN THE TREES: RADIOT	
PREHENSILE TAILED SKINK, CORUCIA ZEBRATA	68
Abstract	69
Introduction	69
Materials and Methods	
Study Area	
Capture and Measurements	
RadiotelemetryDefining Home-Range Size	
Results	72
Movements and Site Fidelity	72
Home-Range Size	
Habitat	
Interactions between Lizards	
Discussion	74
Movements and Site Fidelity	
Home-Range Size	
Habitat	
Interactions between Lizards	
Acknowledgements	
CHAPTER 5: ON THE CENSUS AND EFFECTIVE SIZE	OF A CORUCIA ZERRATA
POPULATION WITH INFERENCES ON THE CONSERVA	
Introduction	

Demographic estimates of N <sub>e</sub>	
N <sub>e</sub> estimated from genetic data	
The census size	
The N <sub>e</sub> /N <sub>c</sub> ratio	
Estimating N <sub>e</sub> in a population of C. zebrata	81
Methods	
Study site, DNA extraction and PCR amplification	
Census N	
Estimation of Ne	83
Results	84
Population census	
Effective population size	
Ne/Nc ratio and rate of genetic drift	85
Discussion	86
Conclusions	88
CHAPTER 6: GENERAL SUMMARY; IMPLICATIONS FOR CONSERVATION MANAGEMENT AND AVENUES OF FURTHER RESEARCH	90
MANAGEMENT AND AVENUES OF FURTHER RESEARCH	09
State of the art before the current study	90
State of the art after the current study	90
PhylogeographyPhylogeography	90
Corucia and social behaviour in the Egernia group	91
Genetic structure, N <sub>c</sub> and N <sub>e</sub>	
Habitat use, home range size and detection probability	93
Future directions	
Improved resolution of colonisation pattern and a molecular test of the sub-species hypothesis	
Social behaviour	
Population vital rates and abundance	
Dispersal and movement pattern	97
Conservation and management	
Conservation status of C. zebrata	_
Conservation in the Solomon Islands and management recommendations for C. zebrata	
Challenges for conservation management in developing countries	100
REFERENCES	102
APPENDIX 1: LIST OF SAMPLES USED IN CHAPTER 2	120
ALLENDIA I. LIOT OF GAMILLEO COLD IN CHAILTEN Z	120
APPENDIX 2: JOURNAL PUBLICATION	123

#### **Abstract**

A complete understanding of species' ecology spans a number of biological disciplines and includes the distribution and genetical lineages and how these lineages correspond with geological features in space and time, the properties of single populations (group composition, neighbourhood size and population size), habitat preferences and the behaviour and movement pattern of single individuals. This study focused on the large and ecologically atypical prehensile-tailed skink (Corucia zebrata), a reptile endemic to the Solomon Archipelago. Due to its large size and unusual ecology, C. zebrata is regarded as one of the most unique reptiles word wide, however very little information about the species exists beyond that of anecdotal nature. This thesis explores the ecology of *C. zebrata* with specific focus on phylogeography, population genetics and population size, habitat preferences and behaviour. Specifically, the aims of the thesis were to 1) Define biogeographic regions for C. zebrata, determine levels of divergence between island populations and assess these patterns in light of information on Pleistocene land bridges, island isolation and island age. 2) Explore the group composition and overall population structure of one C. zebrata population with inferences on social behaviour. 3) Explore the habitat preferences and home range size of C. zebrata with comparisons to other terrestrial and arboreal taxa of similar size. 4) Assess the effective and census population sizes of one C. zebrata population with inferences on the conservation status of the population.

For phylogeographic analysis, 61 *C. zebrata* were sampled from 14 locations across the Solomon Archipelago and sequenced at two mitochondrial loci and four nuclear loci for assessment of population differentiation and Bayesian phylogenetic inference. These data were regarded in light of geological information on Pleistocene land bridges and island age. The fine scale genetic structure of one population was assessed from a sample of 51 lizards captured in a 900 ha study plot within a larger area of continuous rainforest on Ugi Island, Makira Province. Bayesian assignment analysis and pair-wise relatedness between individual lizards based on eight polymorphic DNA microsatellite loci were used to investigate population structure and group composition. Additionally, these loci were used to estimate the effective population size on Ugi Island, and were, together with estimations on population census, used for conservation assessment. For assessment of movement patterns, a six-week study of *C. zebrata* on Ugi Island was conducted to determine home-range sizes and overlap among conspecifics. During this time, 25 lizards were fitted with radio transmitters and were followed for periods of 5–38 days.

Populations of *C. zebrata* on separate islands were found to be genetically distinct from each other, with reciprocal monophyly on mitochondrial DNA. Populations on islands connected by Pleistocene land bridges were marginally less divergent than from other nearby but isolated

islands. There are indications that *C. zebrata* has radiated across the eastern islands of the archipelago within the last 1 - 4 million years. Genetic analysis of the population on Ugi Island indicated that the connectivity within the population was high as all animals constituted a single genetic group and there were indications of a high dispersal rate across the sample area. There was ambiguous evidence for social associations of family groups. Telemetry results indicated that the average home range over the period studied was equivalent to the canopy of one tree. Assessment into population size indicated no immediate concern for the population on Ugi; however this cannot be extrapolated to populations on the remainder of the archipelago.

Data in this thesis suggest that dispersal between islands is very limited and that barriers to gene flow exist within the major islands. Phylogeography of *C. zebrata* is not explained by Pleistocene land bridges. The population genetic results indicated that there may be some short-term family association in this species, but potential social adaptations do not affect the overall genetic structure of the population. It was noted that the conventional survey methods that involve searching tree trunk habitats have low detection probability and that the detection probability is heterogeneous with respect to the habitat (tree species) and region of the country. These are important considerations for further ecological studies of the species, in particular for the purpose of assessing its conservation status.

#### **Publications and Presentations**

The thesis is based on the following articles and conference publications:

#### **Accepted journal publication:**

**Hagen, I.J.** & Bull, C.M. (2011) Home Ranges in the Trees: Radio Telemetry of the Prehensile Tailed Skink, *Corucia zebrata*. *Journal of Herpetology*, **45**, 22-25. This publication appears in Chapter 4.

#### **Submitted journal publication:**

**Hagen**, **I.J.**, Donnellan, S., Herfindal, I. & Bull, C.M. Hanging about in Trees: Population Genetic Structure of the Prehensile Tailed Lizard, *Corucia zebrata*. Submitted to Wildlife Research. This manuscript appears in Chapter 3.

#### **Journal publication awaiting submission:**

**Hagen**, **I.J.**, Donnellan, S. & Bull., C.M. Phylogeography and colonisation history of the prehensile tailed skink *Corucia zebrata* on the Solomon Archipelago. To be submitted to Journal of Biogeography. This manuscript appears in Chapter 2.

#### **Conference publications:**

'Conservation genetics of *Corucia zebrata* and implications for conservation planning and management'. ESF Conservation Genetics Conference, Trondheim, Norway, May 2009. Oral presentation. Data from this presentation appears in Chapter 2.

'Phylogeography of Endemic Taxa Across the Solomon Archipelago'. 6<sup>th</sup> World Congress of Herpetology, Manaus, Brazil, August 2008. Oral presentation. Data from this presentation appears in Chapter 2.

'Genetic N<sub>e</sub> and conservation assessment of elusive species; a case study of the Solomon Islands prehensile tailed skink (*Corucia zebrata*)'. Sustainable Conservation – Bridging the Gap, Trondheim, March 2010. Poster presentation. Data from this presentation appears in Chapter 5.

'Evolution of Social Behaviour in *Egernia* Group of Australasian Skinks'. European Society for Evolutionary Biology (ESEB), Turin, August 2009. Poster presentation. Data from this presentation appears in Chapter 3.

### **Declaration**

I certify that this thesis does not incorporate without acknowledgement any material previously submitted for the degree or diploma in any university; and that to the best of my knowledge and belief it does not contain any material previously published or written by another person except where due reference in made in the text.

INGERID JULIE HAGEN

Ingend julie Hugen

February 2011

This research was carried out under Flinders University

Animal Welfare Committee approval number E230 and a research permit

from the Solomon Island's Department of Education

### Acknowledgement

I would like thank my supervisors, Professors Mike Bull and Steve Donnellan for their help through four years of exciting and sometimes challenging work. During this time my supervisors patiently and generously offered guidance and support.

This work could not have been carried out without the generosity and skilled help of a number of people in the Solomon Islands, Australia and the United States. I would like to thank Mr Michael McCoy for putting me in touch with his friends and family in the Solomon Islands and for providing information about the Solomon Island's culture and the ecology of the study animal and to Dr Chris Filardy for providing advice on the research permit process. I am grateful towards Mrs Janet Wate for providing me with a home whilst spending time in Honiara. Janet and her extended family greatly facilitated the process of establishing myself as a researcher in the Solomon Islands. Mr Patrick Pikacha provided help with field work and logistics and was also excellent company on Makira. Wherever I traveled I was met with generous hospitality: the people of Rendova, Mboremole, Kira Kira, Santa Ana, Kolopakisa, Gizo, Munda, Mt Austen, Choiseul and Shortland Islands overwhelmed me with their kindness. On Ugi Island I would like to extend my gratitude to my hosts Rose and Joseph for their help and hospitality. Emmanuel provided his land on Ugi for my research and helped organizing the climbers. The climbers James and Ray were invaluable; most of the lizards found on Ugi were caught by them. Professor Fred Allendorf and Dr Michael Schwartz at the University of Montana in Missoula generously provided statistical advice. A number of people associated with the South Australian Museum Evolutionary Biology Unit contributed to this work: Annabel Smith and Jacki Smith provided help and excellent company in the field. Luke Price helped with custom made maps and Leanne Wheaton, Kathy Saint, Alison Fitch and Ralph Foster advised in the lab. Dr Adam Skinner provided valuable advice with phylogeographic analysis, whilst Dr Jaro Guzinsky provided statistical advice.

I thank my parents for continuous encouragement throughout the PhD candidature and my partner Ivar for his patience and support in the final and stages of the work.

Lastly, I would like to thank the funding bodies that ultimately made the study possible: National Geographic, Conservation International, the Holsworth Foundation, the Mark Mitchell Foundation, the Environmental Futures Network, the Peter Rankin Trust and the Australian Society of Herpetologists.

This thesis is dedicated to the Rainforest Foundation; a non-governmental organisation that works towards the world wide protection of rainforests.