

6. References

ABAC pol institution of Assumption University n.d., *Evaluation of 30 Baht project*, viewed 1 December 2005, <<http://168.120.31.165/webbase/2546/0630.html>>.

Abramson, JH & Gahlinger, PM 2001, *Computer Programs for Epidemiologists: Papi Version 4.0*, Sagebrush Press, Utah.

Ajayi, AA, Afolabi, MA, Balogun, MO, Adigun, AQ, Ajayi, OE & Akintomide, AO 1996, 'Oral therapy with combined enalapril, prazosin and hydrochlorothiazide in the acute treatment of severe hypertension in Nigerians', *European Journal of Clinical Pharmacology*, vol. 51, no. 1, pp. 45-8.

American Heart Association 2002, *Heart disease and stroke statistics - 2003 update*, viewed 19 October 2003, <<http://www.americanheart.org/downloadable/heart/10590179711482003HDSStatsBookREV7-03.pdf>>.

American Heart Association 2003, *Statistical fact sheet - populations: international cardiovascular disease statistics*, viewed 19 October 2003, <<http://www.americanheart.org/downloadable/heart/1059109452900FS06INT3REV7-03.pdf>>.

Ames, SC, Jones, GN & Brantley, PJ 2001, 'A prospective study of the impact of stress on quality of life: an investigation of low-income individuals with hypertension', *Annals of Behavioral Medicine*, vol. 23, no. 2, pp. 112-119.

Anderson, RB, Nackley, JF & Testa, MA 1995, 'Symptom distress checklists as a component of quality of life measurement: comparing symptom reports with responses to multiple choice questionnaires', *Drug Information Journal*, vol. 29, pp. 1689S-1707S.

Anderson, RB & Testa, MA 1994, 'Symptom distress checklists as a component of quality of life measurement: comparing prompted reports by patients and physician with concurrent adverse event reports via the physician', *Drug Information Journal*, vol. 28, pp. 89-114.

Antonios, TF, Cappuccio, FP, Markandu, ND, Sagnella, GA & MacGregor, GA 1996, 'A diuretic is more effective than a β -blocker in hypertensive patients not controlled on amlodipine and lisinopril', *Hypertension*, vol. 27, no. 6, pp. 1325-1328.

Australian Institute of Health and Welfare 2002, *Australia's health 2002*, Pirie Printers, Canberra.

Australian Institute of Health and Welfare 2004, *The relationship between overweight, obesity and cardiovascular disease*, Australian Institute of Health and Welfare, Canberra.

Baima, J, Nicolaou, M, Schwartz, F, DeStefano, AL, Manolis, A, Gavras, I, Laffer, C, Elijevich, F, Farrer, L, Baldwin, CT & Gavras, H 1999, 'Evidence for linkage between essential hypertension and a putative locus on human chromosome 17', *Hypertension*, vol. 34, no. 1, pp. 4-7.

Bdesha, AS, Bunce, CJ, Snell, ME & Witherow, RO 1994, 'A sham controlled trial of transurethral microwave therapy with subsequent treatment of the control group', *Journal of Urology*, vol. 152, no. 2 Pt 1, pp. 453-458.

Beard, CM, Barnard, RJ, Robbins, DC, Ordavas, JM & Schaefer, EJ 1996, 'Effects of diet and exercise on qualitative and quantitative measures of LDL and its susceptibility to oxidation', *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 16, no. 2, pp. 201-207.

Bender, R, Jockel, KH, Richter, B, Spraul, M & Berger, M 2002, 'Body weight, blood pressure, and mortality in a cohort of obese patients', *American Journal of Epidemiology*, vol. 156, no. 3, pp. 239-245.

Benetos, A, Consoli, S, Safavian, A, Dubanchet, A & Safar, M 2000, 'Efficacy, safety, and effects on quality of life of bisoprolol/hydrochlorothiazide versus amlodipine in elderly patients with systolic hypertension', *American Heart Journal*, vol. 140, no. 4, p. E11.

Bishop, E 2003, 'Reflexology in the management of encopresis and chronic constipation', *Paediatric Nursing*, vol. 15, no. 3, pp. 20-21.

Blais, L, Desgagne, A & LeLorier, J 2000, '3-Hydroxy-3-Methylglutaryl coenzyme a reductase inhibitors and the risk of cancer: a nested case-control study', *Archives of Internal Medicine*, vol. 160, no. 15, pp. 2363-2368.

Brantley, PJ, Jones, GN, Boudreaux, E & Catz, S 1997, 'The weekly stress inventory', in *Evaluating stress: a book of resources*, eds C Zalaquett & R Wood, Scarecrow, Maryland.

Brien, EO, Beevers, DG & Marshall, HJ 1995, *ABC of hypertension*, 3rd edn, BMJ Publishing Group, London.

Broom, A, Barnes, J & Tovey, P 2004, 'Introduction to the research methods in CAM series', *Complementary Therapies in Medicine*, vol. 12, pp. 126-130.

Brown, MJ 1996, 'The causes of essential hypertension', *British Journal of Clinical Pharmacology*, vol. 42, no. 1, July, pp. 21-27.

Brygge, T, Heinig, JH, Collins, P, Ronborg, S, Gehrchen, PM, Hilden, J, Heegaard, S & Poulsen, LK 2001, 'Reflexology and bronchial asthma', *Respiratory Medicine*, vol. 95, no. 3, pp. 173-179.

Byers, DC 2001, *Better health with foot reflexology*, Ingham Publishing Inc., Florida.

Chadwick, DL 1999, 'What are the reasons for nurses using complementary therapy in practice?', *Complementary Therapies in Nursing & Midwifery*, vol. 5, pp. 144-148.

Chaiteerapan, S, Tongtang, V, Puripanyo, K, Sritara, P, Suthichaiyakul, T & Luoatavorn, P 1997, *The research of cardiovascular diseases*, viewed 20 June 2003, <<http://www.thai.net/healthplan/>>.

Chehne, F, Oguogho, A, Lupattelli, G, Palumbo, B & Sinzinger, H 2002, 'Effect of giving up cigarette smoking and restarting in patients with clinically manifested atherosclerosis', *Prostaglandins Leukotrienes & Essential Fatty Acids*, vol. 67, no. 5, pp. 333-339.

Chuprapawan, J, Nasongkla, M, Jirawatkul, A, Kajorntam, Y & Chayowan, N 1995, 'Health interview survey of population aged 50 and over', *Thailand Journal of Health Promotion and Environmental Health*, vol. 20, no. 2, viewed 1 June 2005, <<http://www.anamai.moph.go.th/advisor/202/20201.html>>.

Clark, CC 2000, *Integrating complementary health procedures into practice*, Springer Publishing Company Inc., New York.

Cline, ME, Herman, J, Shaw, E & Morton, RD 1992, 'Standardization of the visual analogue scale', *Nursing Research*, vol. 41, pp. 378-379.

Commonwealth of Australia 1999, *National statement on ethical conduct in research involving humans*, Commonwealth of Australia, Canberra.

Coogan, PF, Rosenberg, L, Palmer, JR, Strom, BL, Zauber, AG & Shapiro, S 2002, 'Statin use and the risk of breast and prostate cancer', *Epidemiology*, vol. 13, no. 3, pp. 262-267.

Cote, I, Moisan, J, Chabot, I & Gregoire, JP 2005, 'Health-related quality of life in hypertension: impact of a pharmacy intervention programme', *Journal of Clinical Pharmacy and Therapeutics*, vol. 30, no. 4, pp. 355-362.

Cyrus-David, MS, Weinberg, A, Thompson, T & Kadmon, D 2005, 'The effect of statins on serum prostate specific antigen levels in a cohort of airline pilots: a preliminary report', *Journal of Urology*, vol. 173, no. 6, pp. 1923-1925.

Dauphinee, S, Gauthier, L, Gandek, B, Magnan, L & Pierre, U 1997, 'Readying a US measure of health status, the SF-36, for use in Canada', *Clinical and Investigative Medicine. Medecine Clinique et Experimentale*, vol. 20, pp. 224-238.

de Jongh, RT, Serne, EH, IJzerman, RG, de Vries, G & Stehouwer, CD 2004, 'Impaired microvascular function in obesity: implications for obesity-associated microangiopathy, hypertension, and insulin resistance', *Circulation*, vol. 109, no. 21, pp. 2529-2535.

Degl'Innocenti, A, Elmfeldt, D, Hofman, A, Lithell, H, Olofsson, B, Skoog, I, Trenkwalder, P, Zanchetti, A & Wiklund, I 2004, 'Health-related quality of life during treatment of elderly patients with hypertension: results from the study on cognition and prognosis in the elderly (SCOPE)', *Journal of Human Hypertension*, vol. 18, no. 4, pp. 239-245.

Devlin, AM, Clark, JS, Reid, JL & Dominiczak, AF 2000, 'DNA synthesis and apoptosis in smooth muscle cells from a model of genetic hypertension', *Hypertension*, vol. 36, no. 1, pp. 110-115.

deWit, SC 1998, *Essentials of medical-surgical nursing*, 4th edn, WB Saunders Company, Pennsylvania.

Dias da Costa, JS, Fuchs, SC, Olinto, MT, Gigante, DP, Menezes, AM, Macedo, S & Gehrke, S 2002, 'Cost-effectiveness of hypertension treatment: a population-based study', *Sao Paulo Medical Journal*, vol. 120, no. 4, pp. 100-104.

Dickerson, LM & Gibson, MV 2005, 'Management of hypertension in older persons', *American Family Physician*, vol. 71, no. 3, pp. 1-10.

Dimenas, E, Dahlof, B & Oloffson, B & Wiklund, I 1990, 'An instrument for quantifying subjective symptoms among untreated hypertensives: development and documentation', *Journal of Clinical Research and Pharmacoepidemiology*, vol. 4, pp. 205-217.

Domenico, GD & Wood, EC 1997, *Beard's massage*, 4th edn, WB Saunders Company, Pennsylvania.

Dominiczak, AF, Negrin, DC, Clark, JS, Brosnan, MJ, McBride, MW & Alexander, MY 2000, 'Genes and hypertension: from gene mapping in experimental models to vascular gene transfer strategies', *Hypertension*, vol. 35, no. 1 part 2, pp. 164-172.

Dougans, I 2002, *Complete illustrated guide to reflexology*, Harper Collins Publishers, London.

Duncan, LE, Lewis, C, Smith, CE, Jenkins, P, Nichols, M & Pearson, TA 2001, 'Sex, drugs, and hypertension: a methodological approach for studying a sensitive subject', *International Journal of Impotence Research*, vol. 13, no. 1, pp. 31-40.

DuPen, S & DuPen, A 2000, *Opioid conversion calculator*, computer software, Cynergy group, Poulsbo, Washington.

Dupuy, HJ 1984, 'The psychological general well-being (PGWB) index', in *Assessment of quality of life in clinical trials of cardiovascular therapies*, eds NK Wenger, ME Mattson, CD Furberg & J Elinson, Le Jacq publishing, New York.

Elliott, P 1991, 'Observational studies of salt and blood pressure', *Hypertension*, vol. 17, pp. 13-18.

Erickson, SR, Williams, BC & Gruppen, LD 2001, 'Perceived symptoms and health-related quality of life reported by uncomplicated hypertensive patients compared to normal controls', *Journal of Human Hypertension*, vol. 15, pp. 539-548.

Erickson, SR, Williams, BC & Gruppen, LD 2004, 'Relationship between symptoms and health-related quality of life in patients treated for hypertension', *Pharmacotherapy*, vol. 24, no. 3, pp. 344-350.

Evans, SL, Nokes, LDM, Weaver, P, Maheson, M & Morrell, P 1998, 'Effect of reflexology treatment on recovery after total knee replacement', *Journal of Bone & Joint Surgery*, vol. 80, p. 172.

Faergeman, O & Grundy, S 2003, *Dyslipidaemia*, Elsevier Science Ltd, London.

Farnsworth, P 1995, *The Australian College of Tactile Therapies: Reflexology Seminar*, The Australian College of Tactile Therapies, Adelaide, Australia.

Feeny, D, Furlong, W, Boyle, M & Torrance, GW 1995, 'Multi-attribute health status classification systems: health utilities index', *Pharmacoeconomics*, vol. 7, pp. 490-502.

Feigin, V, Parag, V, Lawes, CM, Rodgers, A, Suh, I, Woodward, M, Jamrozik, K & Ueshima, H 2005, 'Smoking and elevated blood pressure are the most important risk factors for subarachnoid hemorrhage in the Asia-Pacific region', *Stroke*, vol. 36, no. 7, pp. 1360-1365.

Ferreira, MI, Alcocer, L, Leary, WP & Reyes, AJ 1992, 'Quality of life during antihypertensive therapy with diuretics: critical reappraisal', in AJ Reyes (ed), *Progress in pharmacology and clinical pharmacology, vol. 9, Diuretics: clinical pharmacology and uses in cardiovascular medicine, nephrology and hepatology*, Fischer, Stuttgart.

Ferrell-Torry, AT & Glick, OJ 1993, 'The use of therapeutic massage as a nursing intervention to modify anxiety and the perception of cancer pain', *Cancer Nursing*, vol. 16, no. 2, pp. 93-101.

Fogari, R & Zoppi, A 2002, 'Effects of antihypertensive therapy on sexual activity in hypertensive men', *Current Hypertension Reports*, vol. 4, pp. 202-210.

Frankel, BSM 1997, 'The effects of reflexology on baroreceptor reflex sensitivity, blood pressure and sinus arrhythmia', *Complementary Therapies in Medicine*, vol. 5, pp. 80-84.

Franz, MH, Schmieder, RE & Matthew, W 1997, 'Salt: a perpetrator of hypertensive target organ disease?', *Archives of Internal Medicine*, vol. 157, no. 21, pp. 2449-2452.

Fraser, J & Kerr, JR 1993, 'Psychophysiological effects of back massage on elderly institutionalized patients', *Journal of Advanced Nursing*, vol. 18, pp. 238-245.

Frethiem, A. 2003, 'Back to thiazide-diuretics for hypertension: reflections after a decade of irrational prescribing', *BMC Family Practice*, vol. 4, viewed 28 August 2005, (online OVID/Journals@OVID).

Fretheim, A, Aaserud, M & Oxman, AD 2003, 'The potential savings of using thiazides as the first choice antihypertensive drug: cost-minimisation analysis', *BMC Health Services Research*, vol. 3, viewed 28 August 2005, (online OVID/Journals@OVID).

Friedman, R, Schwartz, JE, Schnall, PL, Landsbergis, PA, Pieper, C, Gerin, W & Pickering, TG 2001, 'Psychological variables in hypertension: relationship to casual or ambulatory blood pressure in men', *Psychosomatic Medicine*, vol. 63, no. 1, pp. 19-31.

Fuchs, FD, Chambless, LE, Whelton, PK, Nieto, FJ & Heiss, G 2001, 'Alcohol consumption and the incidence of hypertension: the atherosclerosis risk in communities study', *Hypertension*, vol. 37, no. 5, pp. 1242-1250.

Gambles, M, Crooke, M & Wilkinson, S 2002, 'Evaluation of a hospice based reflexology service: a qualitative audit of patient perceptions', *European Journal of Oncology Nursing*, vol. 6, pp. 37-44.

Georgiades, A, Sherwood, A, Gullette, EC, Babvak, MA, Hinderliter, A, Waugh, R, Tweedy, D, Craighead, L, Bloomer, R & Blumenthal, JA 2000, 'Effects of exercise

and weight loss on mental stress-induced cardiovascular responses in individuals with high blood pressure', *Hypertension*, vol. 36, no. 2, pp. 171-176.

Glynn, RJ, Brock, DB, Harris, T, Havlik, RJ, Chrischilles, EA, Ostfeld, AM, Taylor, JO & Hennekens, CH 1995, 'Use of antihypertensive drugs and trends in blood pressure in the elderly', *Archives of Internal Medicine*, vol. 155, no. 17, pp. 1855-1860.

Gottdiener, JS, Reda, DJ, Massie, BM, Materson, BJ, Williams, DW & Anderson, RJ 1997, 'Effect of single-drug therapy on reduction of left ventricular mass in mild to moderate hypertension: comparison of six antihypertensive agents', *Circulation*, vol. 95, no. 8, pp. 2007-2014.

Grealish, L, Lomasney, A & Whiteman, B 2000, 'Foot massage: a nursing intervention to modify the distressing symptoms of pain and nausea in patients hospitalized with cancer', *Cancer Nursing*, vol. 23, pp. 237-243.

Gress, TW, Nieto, FJ, Shahar, E, Wofford, MR & Brancati, FL 2000, 'Hypertension and antihypertensive therapy as risk factors for type 2 diabetes mellitus', *New England Journal of Medicine*, vol. 342, pp. 905-912.

Groppelli, A, Giorgi, DM, Omboni, S, Parati, G & Mancia, G 1992, 'Persistent blood pressure increase induced by heavy smoking', *Journal of Hypertension*, vol. 10, no. 5, pp. 495-499.

Gueyffier, F, Boutitie, F, Boissel, JP, Pocock, S, Coope, J, Cutler, J, Ekblom, T, Fagard, R, Friedman, L, Perry, M, Prineas, R & Schron, E 1997, 'Effect of antihypertensive drug treatment on cardiovascular outcomes in women and men', *Annals of Internal Medicine*, vol. 126, no. 10, pp. 761-767.

Guyatt, GH, Feeny, DH & Patrick, DL 1993, 'Measuring Health-related Quality of Life', *Annals of Internal Medicine*, volume 118, no. 8, pp. 622-629.

Hajjar, IM, Grim, CE, George, V & Kotchen, TA 2001, 'Impact of diet on blood pressure and age-related changes in blood pressure in the US population: analysis of NHANES III', *Archives of Internal Medicine*' vol. 161, no. 4, pp. 589-593.

Halbert, JA, Silagy, CA, Finucane, P, Withers, RT, Hamdorf, PA & Andrews, GR 1997, 'The effectiveness of exercise training in lowering blood pressure: a meta-analysis of randomized controlled trials of 4 weeks or longer', *Journal of Human Hypertension*, vol. 11, no. 10, pp. 641-649.

Halimi, JM, Giraudeau, B, Vol, S, Caces, E, Nivet, H & Tichet, J et al. 2002, 'The risk of hypertension in men: direct and indirect effects of chronic smoking', *Journal of Hypertension*, vol. 20, no. 2, pp. 187-193.

Harangi, M, Seres, I, Varga, Z, Emri, G, Szilvassy, Z, Paragh, G & Remenik, E 2004, 'Atorvastatin effect on high-density lipoprotein-associated paraoxonase activity and oxidative DNA damage', *European Journal of Clinical Pharmacology*, vol. 60, no. 10, pp. 685-691.

Harrington, F, Saxby, BK, McKeith, IG, Wesnes, K & Ford, GA 2000, 'Cognitive performance in hypertensive and normotensive older subjects', *Hypertension*, vol. 36, no. 6, pp. 1079-1082.

Hattan, J, King, L & Griffiths, P 2002, 'The impact of foot massage and guided relaxation following cardiac surgery: a randomized controlled trial', *Journal of Advanced Nursing*, vol. 37, pp. 199-207.

Hayashi, T, Boyko, EJ, Leonetti, DL, McNeely, MJ, Newell-Morris, L, Kahn, SE & Fujimoto, WY 2004, 'Visceral adiposity is an independent predictor of incident hypertension in Japanese Americans', *Annals of Internal Medicine*, vol. 140, no. 12, pp. 992-1000.

Hayes, J & Cox, C 1999, 'Immediate effects of a five-minute foot massage on patients in critical care', *Intensive and Critical Care Nursing*, vol. 15, pp. 83-94.

Hays, RD, Sherbourne, CD & Mazel, RM 1993, 'The rand 36-item health survey 1.0', *Health Economy*, vol. 2, pp. 217-227.

He, FJ & MacGregor, GA 2003, 'How far should salt intake be reduced?', *Hypertension*, vol. 42, pp. 1093-1099.

He, FJ, Markandu, ND & MacGregor, GA 2005, 'Modest salt reduction lowers blood pressure in isolated systolic hypertension and combined hypertension', *Hypertension*, vol. 46, pp. 66-70.

He, J, Muntner, P, Chen, J, Roccella, EJ, Streiffer, RH & Whelton, PK 2002, 'Factors associated with hypertension control in the general population of the United States', *Archives of Internal Medicine*, vol. 162, no. 9, pp. 1051-1058.

He, J, Whelton, PK, Appel, LJ, Charleston, J & Klag, MJ 2000, 'Long-term effects of weight loss and dietary sodium reduction on incidence of hypertension', *Hypertension*, vol. 35, no. 2, pp. 544-549.

Healey, J 2002, *Stress*, The Spinney Press, NSW, Australia.

Helmets, KF, Baker, B, O'Kelly, B & Tobe, S 2000, 'Anger expression, gender, and ambulatory blood pressure in mild, unmedicated adults with hypertension', *Annals of Behavioral Medicine*, vol. 22, no. 1, pp. 60-64.

Hernandez-Reif, M, Field, T, Field, T & Theakston, H 1998, 'Multiple sclerosis patients benefit from massage therapy', *Journal of Bodywork and Movement Therapies*, vol. 3, no. 3, pp. 168-174.

Hodgson, H 2000, 'Does reflexology impact on cancer patients' quality of life?', *Nursing Standard*, vol. 14, no. 31, pp. 33-38.

Hoes, AW, Grobbee, DE, Lubsen, J, Man in't Veld, AJ, van der Does, E & Hofman, A 1995, 'Diuretics, beta-blockers, and the risk for sudden cardiac death in hypertensive patients', *Annals of Internal Medicine*, vol. 123, no. 7, pp. 481-487.

Holmes, S & Dickerson, J 1987, 'The quality of life: design and evaluation of a self-assessment instrument for use in cancer patients', *International Journal of Nursing Studies*, vol. 24, pp. 15-24.

Hulme, J, Waterman, H & Hillier, VF 1999, 'The effect of foot massage on patients' perception of care following laparoscopic sterilization as day case patients', *Journal of Advanced Nursing*, vol. 30, pp. 460-468.

Hunter, M 1992, 'The women's health questionnaire: a measure of mid-aged women's perceptions of their emotional and physical health', *Psychology and Health*, vol. 7, pp. 45-54.

Hunter, SJ, Wiggam, MI, Ennis, CN, Whitehead, HM, Sheridan, B, Atkinson, AB & Bell, PM 1999, 'Comparison of effects of captopril used either alone or in combination with a thiazide diuretic on insulin action in hypertensive type 2 diabetic patients: a double-blind crossover study', *Diabetic Medicine*, vol. 16, no. 6, pp. 482-487.

Ibrahim, J, Schachter, M, Hughes, AD & Sever, PS 1999, 'Low dose indapamide plus perindopril combination effects on cardiovascular structure and function in genetic hypertension', *Clinical and Experimental Pharmacology and Physiology*, vol. 26, pp. 622-627.

Imamura, H, Katagiri, S, Uchid, K, Miyamoto, N, Nakano, H & Shirota, T 2000, 'Acute effects of moderate exercise on serum lipids, lipoproteins and apolipoproteins

in sedentary young women', *Clinical and Experimental Pharmacology and Physiology*, vol. 27, no. 12, pp. 975-979.

Ishizaka, N, Ishizaka, Y, Toda, E, Hashimoto, H, Nagai, R & Yamakado, M 2005, 'Association between cigarette smoking, metabolic syndrome, and carotid arteriosclerosis in Japanese individuals', *Atherosclerosis*, vol. 181, no. 2, pp. 381-388.

Ismail, MK 2005, *Diet for coronary heart disease and artherosclerosis - part I: what causes coronary heart disease?*, viewed 28 October 2005, <<http://www.bawarchi.com/health/coronaryheart-diet1.html>>.

Janzon, E, Hedblad, B, Berglund, G & Engstrom, G 2004, 'Changes in blood pressure and body weight following smoking cessation in women', *Journal of Internal Medicine*, vol. 255, pp. 266-272.

Jia, H, Hingorani, AD, Sharma, P, Hopper, R, Dickerson, C, Trutwein, D, Lloyd, DD & Brown, MJ 1999, 'Association of the G_sα gene with essential hypertension and response to β-blockade', *Hypertension*, vol. 34, no. 1, pp. 8-14.

Jirayingmongkol, P, Chantein, S, Phengchomjan, N & Bhanggananda, N 2002, 'The effect of foot massage with biofeedback: a pilot study to enhance health promotion', *Nursing and Health Sciences*, vol. 4, no. 3 Suppl, p. A4.

Joint Commission on the Accreditation of Healthcare Organizations 2001, 'Pain standards for 2001: an exploratory study of reflexological treatment for headache', *Alternative Therapies in Health and Medicine*, vol. 5, pp. 57-65.

Joint National Committee 2003, *Prevention, detection, evaluation and treatment of high blood pressure*, seventh report (AV Chobanian, chairman), The National Institutes of Health, Maryland

Jovelic, S, Hajdukovic, Z, Jovelic, A & Raden, S 2005, 'Cigarette smoking in military pilots and intima-media thickness of the carotid arteries', *Vojnosanitetski Pregled*, vol. 62, no. 5, pp. 365-370.

Kalimo, R., & Vuori, J 1993, 'Psychosocial factors in youth and at working age as predictors of blood pressure: a prospective study', *Journal of Psychosomatic Research*, vol. 37, no. 6, Sept, pp. 643-652.

Kaplan, NM, Lieberman, E & Neal, W 2002, *Kaplan's clinical hypertension*, 8th edn, Lippincott Williams & Wilkins, Philadelphia.

Keegan, L 2001, *Healing with complementary & alternative therapies*, Delmar, New York.

Khoury, S, Yarows, SA, O'Brien, TK & Sowers, JR 1992, 'Ambulatory blood pressure monitoring in a non-academic setting: effects of age and sex', *American Journal of Hypertension*, vol. 5, no. 9, pp. 616-623.

Kinjo, K, Sato, H, Sakata, Y, Nakatani, D, Mizuno, H, Shimizu, M, Sasaki, T, Kijima, Y, Nishino, M, Uematsu, M, Tanouchi, J, Nanto, S, Otsu, K, Hori, M & Osaka 2005, 'Impact of smoking status on long-term mortality in patients with acute myocardial infarction', *Circulation Journal*, vol. 69, no. 1, pp. 7-12.

Kohara, H, Miyauchi, T, Suehiro, Y, Ueoka, H, Takeyama, H & Morita, T 2004, 'Combined modality treatment of aromatherapy, footsoak, and reflexology relieves fatigue in patients with cancer', *Journal of Palliative Medicine*, vol. 7, no. 6, pp. 791-796.

Kotsis, V, Stabouli, S, Bouldin, M, Low, A, Toumanidis, S & Zakopoulos, N 2005, 'Impact of obesity on 24-hour ambulatory blood pressure and hypertension', *Hypertension*, vol. 45, pp. 602-607.

Kowalak, JP, Chohan, ND & Follin, SA 2003, *Nurse's handbook of alternative & complementary therapies*, Lippincott Williams & Wilkins, Philadelphia.

Kraus, WE, Houmard, JA, Duscha, BD, Knetzger, KJ, Wharton, MB, McCartney, JS, Bales, CW, Henes, S, Samsa, GP, Otvos, JD, Kulkarni, KR & Slentz, CA 2002, 'Effects of the amount and intensity of exercise on plasma lipoproteins', *New England Journal of Medicine*, vol. 347, no. 19, pp. 1483-1492.

Kris-Etherton, PM, Pearson, TA, Wan, Y, Hargrove, RL, Moriarty, K & Fishell, V 1999, 'High-monosaturated fatty acid diets lower both plasma cholesterol and

triacylglycerol concentrations', *American Journal of Clinical Nutrition*, vol. 70, no. 6, pp. 1009-1015.

Kronig, B, Pittrow, DB, Kirch, W, Welzel, D & Weidinger, G 1997, 'Different concepts in first-line treatment of essential hypertension: comparison of a low-dose reserpine-thiazide combination with nitrendipine monotherapy', *Hypertension*, vol. 29, no. 2, pp. 651-658.

Kuhn, MA 1999, *Complementary therapies for health care providers*, Lippincott Williams & Wilkins, Philadelphia.

Kunz, B & Kunz, K 1999, *The complete guide to foot reflexology*, 2nd edn, Harper Collins, London.

Kushiro, T, Takahashi, A, Saito, F, Otsuka, Y, Soma, M, Kurihara, T, Satomura, A, Saito, T & Kanmatsuse, K 2005, 'Erectile dysfunction and its influence on quality of life in patients with essential hypertension', *American Journal of Hypertension*, vol. 18, no. 3, pp. 427-430.

Lacourciere, Y, Belanger, A, Godin, C, Halle, JP, Ross, S, Wright, N & Marion, J 2000, 'Long term comparison of losartan and enalapril on kidney function in hypertensive type 2 diabetics with early nephropathy', *Kidney International*, vol. 58, no. 2, pp. 762-769.

Lalonde, L, O'Connor, A, Joseph, L & Grover, SA 2004, 'Health-related quality of life in cardiac patients with dyslipidemia and hypertension', *Quality of Life Research*, vol. 13, no. 4, pp. 793-804.

Launso, L, Brendstrup, E & Arnberg, S 1999, 'An exploratory study of reflexological treatment for headache', *Alternative Therapies in Health and Medicine*, vol. 5, no. 3, pp. 57-65.

Le Fevre, P, Devereux, J, Smith, S, Lawrie, SM & Cornbleet, M 1999, 'Screening for psychiatric illness in the palliative care inpatient setting: a comparison between the hospital anxiety and depression scale and the general health questionnaire', *Palliative Medicine*, vol. 13, pp. 399-407.

Lea, JP & Nicholas, SB 2002, 'Diabetes and hypertension: key risk factors for kidney disease', *Journal of the National Medical Association*, vol. 94, no. 8, pp. 7s – 15s.

Lee, DH, Ha, MH, Kim, JR & Jacobs, Dr Jr 2001, 'Effects of smoking cessation on changes in blood pressure and incidence of hypertension: a 4-year follow-up study', *Hypertension*, vol. 37, no. 2, pp. 194-198.

Levenstein, S, Smith, MW & Kaplan, GA 2001, 'Psychosocial predictors of hypertension in men and women', *Archives of Internal Medicine*, vol. 161, no. 10, pp. 1341-1346.

Liang, YL, Shiel, LM, Teede, H, Kotsopoulos, D, McNeil, J, Cameron, JD & McGrath, BP 2001, 'Effects of blood pressure, smoking, and their interaction on carotid artery structure and function', *Hypertension*, vol. 37, pp. 6-11.

Long, L, Huntley, A & Ernst, E 2001, 'Which complementary and alternative therapies benefit which conditions? A survey of the opinions of 223 professional organizations', *Complementary Therapies in Medicine*, vol. 9, pp. 178-185.

Mackereth, PA & Tiran, D 2002, *Clinical reflexology: a guide for health professionals*, Churchill Livingstone, London.

MacMahon, S, Peto, R, Cutler, J, Collins, R, Sorlie, P, Neaton, J, Abbott, R, Godwin, J, Dyer, A & Stamler, J 1990, 'Blood pressure, stroke, and coronary heart disease: part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias', *Lancet*, vol. 335, no. 8692, pp. 765-774.

Malinski, MK, Sesso, HD, Lopez-Jimenez, F, Buring, JE & Gaziano, JM 2004, 'Alcohol consumption and cardiovascular disease mortality in hypertensive men', *Archives of Internal Medicine*, vol. 164, no. 6, pp. 623-628.

Mancia, G, Julius, S, Saruta, T, Weber, MA, Ferrari, AU & Wilkinson, IB 2002, *Manual of hypertension*, Harcourt Publishers Ltd, London.

Manger, WM & Gifford, RW 2001, *100 questions and answers about hypertension*, Blackwell Science Inc., Massachusetts.

Mansur, AP, Serrano, CV Jr, Nicolau, JC, Cesar, LA & Ramires, JA 1999, 'Effect of cholesterol lowering treatment on positive exercise tests in patients with hypercholesterolaemia and normal coronary angiograms', *Heart*, vol. 82, no. 6, pp. 689-693.

Maris, ME, Melchert, RB, Joseph, J & Kennedy, RH 2005, 'Gender differences in blood pressure and heart rate in spontaneously hypertensive and wistar-kyoto rats', *Clinical and Experimental Pharmacology and Physiology*, vol. 32, pp. 35-39.

Marques-Vidal, P, Arveiler, D, Evans, A, Amouvel, P, Ferrieres, J & Ducimetiere, P 2001, 'Different alcohol drinking and blood pressure relationships in France and Northern Ireland: the PRIME study', *Hypertension*, vol. 38, no. 6, pp. 1361-1366.

Mathers, C & Penm, R 1999, *Health system costs of cardiovascular diseases and diabetes in Australia 1993-94*, Australian Institute of Health and Welfare, Canberra.

Matthews, KA, Kiefe, CI, Lewis, CE, Liu, K, Sidney, S & Yunis, C 2002, 'Socioeconomic trajectories and incident hypertension in a biracial cohort of young adults', *Hypertension*, vol. 39, no. 3, pp. 772-776.

McCabe, P 2001, *Complementary therapies in nursing and midwifery: from vision to practice*, Ausmed publications, Melbourne.

McCaffery, M & Pasero, C 1999, *Pain: clinical manual*, 2nd edn, Mosby, St. Louis, Missouri.

McDowell, I & Newell, C 1996, *Measuring Health: a guide to rating scales and questionnaires*, 2nd edn, Oxford university press, Inc., New York.

McGuire, DB 1988, 'Measuring pain', in *Instruments for clinical nursing research*, ed M Frank-Stromborg, Appleton an Lange, Norwalk, Connecticut.

McKenney, JM 2005, 'Efficacy and safety of rosuvastatin in treatment of dyslipidemia', *American Journal of Health System Pharmacy*, vol. 62, pp. 1033-1047.

Melzack, R 1987, 'The Short-Form McGill Pain Questionnaire: major properties and scoring methods', *Pain*, vol. 30, pp. 191-197.

Mendez, MA, Cooper, R, Wilks, R, Luke, A & Forrester, T 2003, 'Income, education, and blood pressure in adults in Jamaica, a middle-income developing country', *International Journal of Epidemiology*, vol. 32, no. 3, pp. 400-408.

Midgley, JP, Matthew, AG, Greenwood, CM & Logan, AG 1996, 'Effect of reduced dietary sodium on blood pressure: a meta-analysis of randomized controlled trials', *Journal of the American Medical Association*, vol. 275, no. 20, pp. 1590-1597.

Mikkelsen, KL, Wiinberg, N, Hoegholm, A, Christensen, HR, Bang, LE, Nielsen, PE, Svendsen, TL, Kampmann, JP, Madsen, NH & Bentzon, MW 1997, 'Smoking related to 24-h ambulatory blood pressure and heart rate', *American Journal of Hypertension*, vol. 10, no. 5 Pt 1, pp. 483-491.

Milligan, M, Fanning, M, Hunter, S, Tadjali, M & Stevens, E 2002, 'Reflexology audit: patient satisfaction, impact on quality of life and availability in Scottish hospices', *International Journal of Palliative Nursing*, vol. 8, no. 10, pp. 489-496.

Ministry of Public Health 2001, *Public health statistics A.D. 2001*, Ministry of Public Health, Bangkok, Thailand.

Monahan, FD & Neighbors, M 1998, *Medical-surgical nursing foundations for clinical practice*, 2nd edn, WB Saunders Company, Pennsylvania.

Morris, P, Raphael, B & Bordujenko, A 1999, *Stress and challenge: health and disease*, Commonwealth of Australia, Brisbane.

Mosler, Al, Mignone, AM, Reyes, AJ, Olhaberry, JV, Chiesa, PD, Batista, LB & Santucci, MR 1993, 'Semi-personalized evaluation of quality of life in a study of 25 mg hydrochlorothiazide and a non-diuretic dose of toseamide once daily in elderly hypertensives', in JB Pruschett, A Greenberg (eds), *Diuretics IV: chemistry, pharmacology and clinical applications, internal congress series, vol. 1023*, Elsevier Science Publishers, Amsterdam.

Nakanishi, N 2005, 'Daily life activity and the risk of developing hypertension in middle-aged Japanese men', *Archives of Internal Medicine*, vol. 165, pp. 214-220.

Nanchadal, K, Ashton, D & Wood, DA 2000, 'Alcohol consumption, metabolic cardiovascular risk factors and hypertension in women', *International Journal of Epidemiology*, vol. 29, pp. 57-64.

Naruse, Y, Nakagawa, H, Kagamimori, S, Fujita, Y, Hashimoto, T, Kasamatsu, T, Mikawa, K & Ueshima, H 1998, 'Relationship of parental history of high blood pressure to blood pressure: combined findings of three Japanese population samples, the INTERSALT study', *Journal of Human Hypertension*, vol. 12, no. 4, pp. 215-220.

National Economic and Social Development Board 1997, *Public Health Development Plan in National Economic & Social Development Plan Version 8 (1997-2001)*, National Economic and Social Development Board, Bangkok, Thailand.

National Heart Foundation of Australia 2003, *Hypertension management guide for doctors 2004*, National Heart Foundation of Australia, Canberra.

National Library of Thailand Cataloguing in Publication Data 2001, *Thailand public health 2001*, 3rd edn, Alpha Research Co. Ltd, Bangkok, Thailand.

Neter, JE, Stam, BE, Kok, FJ, Grobbee, DE & Geleijnse, JM 2003, 'Influence of weight reduction on blood pressure: a meta-analysis of randomised controlled trials', *Hypertension*, vol. 42, no. 5, pp. 878-884.

Newman, TB & Hulley, SB 1996, 'Carcinogenicity of lipid-lowering drugs', *The Journal of the American Medical Association*, vol. 275, pp. 55-60.

Nilsson, P, Ostergren, PO, Lindholm, L & Schersten, B 1994, 'Can social class differentials in hypertension be explained by the general susceptibility hypothesis?', *Social Science & Medicine*, vol. 38, no. 9, pp. 1235-1242.

Niskanen, L, Laaksonen, DE, Nivvsson, K, Punnonen, K, Valkonen, VP, Fuentes, R, Tuomainen, TP, Salonen, R & Salonen, JT 2004, 'Inflammation, abdominal obesity, and smoking as predictors of hypertension', *Hypertension*, vol. 44, no. 6, pp. 859-865.

Okuyama, T, Akecho, T, Kugaya, A, Okamura, H, Shima, Y, Maruguchi, M, Hosaka, T & Uchitomi, Y 2000, 'Development and validation of the cancer fatigue scale: a brief, three-dimensional, self-rating scale for assessment of fatigue in cancer patients', *Journal of Pain and Symptom management*, vol. 19, pp. 5-14.

Oleson, T & Flocco, W 1993, 'Randomized controlled study of premenstrual symptoms treated with ear, hand, and foot reflexology', *Obstetrics and Gynaecology*, vol. 82, pp. 906-911.

Pan, W, Pintar, T, Anton, J, Lee, VV, Vaughn, WK & Collard, CD 2004, 'Statins are associated with a reduced incidence of perioperative mortality after coronary artery bypass graft surgery', *Circulation*, vol. 110, no. 11 suppl 1, pp. II45-II49).

Panagiotakos, DB, Pitsavos, C, Chrysohoou, C, Skoumas, J, Zeimbekis, A, Papaioannou, I & Stefanadis, C 2003, 'Effect of leisure time physical activity on blood lipid levels: the ATTICA study', *Coronary Artery Disease*, vol. 14, no. 8, pp. 533-539.

Park, HS & Cho, GY 2004, 'Effects of foot reflexology on essential hypertension patients', *Daehan Ganho Haghoeji*, vol. 34, no. 5, pp. 739-750.

Pescatello, LS, Murphy, D & Costanzo, D 2000, 'Low-intensity physical activity benefits blood lipids and lipoproteins in older adults living at home', *Age and Ageing*, vol. 29, pp. 433-439.

Pickering, TG, Devereux, RB, James, GD, Gerin, W, Landsberdis, P, Schnall, PL & Schwartz, JE 1996, 'Environmental influences on blood pressure and the role of job strain', *Journal of Hypertension – Supplement*, vol. 14, no. 5, pp. S179-S185.

Poirier, P, Lemieux, I, Mauriege, P, Dewailly, E, Blanchet, C, Bergeron, J & Despres, JP 2005, 'Impact of waist circumference on the relationship between blood pressure and insulin: the Quebec health survey', *Hypertension*, vol. 45, no.3, pp. 363-367.

Pourmand, G, Alidaee, MR, Rasuli, S, Maleko, A & Mehrsai, A 2004, 'Do cigarette smokers with erectile dysfunction benefit from stopping?', *BJU International*, vol. 94, no. 9, pp. 1310-1313.

Primatesta, P, Falaschetti, E, Gupta, S, Marmot, MG & Poulter, NR 2001, 'Association between smoking and blood pressure: evidence from the health survey for England', *Hypertension*, vol. 37, no. 2, pp. 187-193.

Rankin-Box, D 2001, *The nurse's handbook of complementary therapies*, Harcourt Publishers Ltd, London.

Renaud, SC, Gueguen, R, Conard, P, Lanzmann-Petithory, D, Orgogozo, JM & Henry, O 2004, 'Moderate wine drinkers have lower hypertension-related mortality: a prospective cohort study in French men', *American Journal of Clinical Nutrition*, vol. 80, no. 3, pp. 621-625.

Renwick, R, Brown, I & Nagler, M 1996, *Quality of life in health promotion and rehabilitation: conceptual approaches, issues, and applications*, SAGE Publications Inc., California.

Reynolds, K, Lewis, B, Nolen, JD, Kinney, GL, Sathya, B & He, J 2003, 'Alcohol consumption and risk of stroke: a meta-analysis', *Journal of the American Medical Association*, vol. 289, no. 5, pp. 579-588.

Richards, KC 1998, 'Effect of a back massage and relaxation intervention on sleep in critical ill patients', *American Journal of Critical Care*, vol. 7, no. 4, pp. 288-299.

Rosen, RC, Cappelleri, JC & Gendrano, N 2002, 'The international index of erectile function (IIEF): a state-of-the-science review', *International Journal of Impotence Research*, vol. 14, pp. 226-244.

Ross, CS, Hamilton, J, Macrae, G, Docherty, C, Gould, A & Cornbleet, MA 2002, 'A pilot study to evaluate the effect of reflexology on mood and symptom rating of advanced cancer patients', *Palliative Medicine*, vol. 16, no. 6, pp. 544-545.

Royal College of Nursing, Australia 1997, *Complementary therapies in Australian nursing practice*, viewed 28 November 2005, <http://www.rcna.org.au/content/complementary_therapies_march_03_-_under_review_25nov04.pdf>.

Rutledge, T & Linden, W 2003, 'Defensiveness and 3-year blood pressure levels among young adults: the mediating effect of stress-reactivity', *Annals of Behavioral Medicine*, vol. 25, no. 1, pp. 34-40.

Sanders, MW, Fazzi, GE, Janssen, GM, Blanco, CE & De Mey, JG 2005, 'High sodium intake increases blood pressure and alters renal function in intrauterine growth-retarded rats', *Hypertension*, vol. 46, no. 1, pp. 71-75.

Sarason, IG, Johnson, JH & Siegel, JM 1978, 'Assessing the impact of life changes: development of the life experiences survey', *Journal of Consulting and Clinical Psychology*, vol. 46, pp. 932-946.

Sareli, P, Radevski, IV, Valtchanova, ZP, Libhaber, E, Candy, GP, DenHond, E, Libhaber, C, Skudicky, D, Wang, JG, Staessen, JA 2001, 'Efficacy of different drug classes used to initiate antihypertensive treatment in black subjects: results of a randomized trial in Johannesburg, South Africa', *Archives of Internal Medicine*, vol. 161, no. 7, pp. 965-971.

Saxby, BK, Harrington, F, McKeith, IG, Wesnes, K & Ford, GA, 2003, 'Effects of hypertension on attention, memory, and executive function in older adults', *Health Psychology*, vol. 22, no. 6, pp. 587-591.

Schalock, RL 1996, *Quality of life volume 1: conceptualization and measurement*, American Association on Mental Retardation, Washington.

Schminke, U, Luedemann, J, Berger, K, Alte, D, Mitusch, R, Wood, WG, Jaschinski, A, Barnow, S, John, U & Kessler, C 2005, 'Association between alcohol consumption and subclinical carotid atherosclerosis: the study of health in Pomerania', *Stroke*, vol. 36, no. 8, pp. 1746-1752.

Schnall, PL, Pieper, C, Schwartz, JE, Karasek, RA, Schlusser, Y, Devereux, RB, Ganau, A, Alderman, M, Warren, K & Pickering, TG 1990, 'The relationship between 'job strain', workplace diastolic blood pressure, and left ventricular mass index: results of a case-control study', *Journal of the American Medical Association*, vol. 263, no. 14, pp. 1929-1935.

Seaward, BL 1999, *Managing stress: principles and strategies for health and wellbeing*, Jones and Barlett Publishers International, London.

Shahinfar, S, Simpson, RL, Carides, AD, Thiyagarajan, B, Nakagawa, Y, Umans, JG, Parks, JH & Coe, FL 1999, 'Safety of losartan in hypertensive patients with thiazide-induced hyperuricemia', *Kidney International*, vol. 56, no. 5, pp. 1879-1885.

Shaw, WS, Patterson, TL, Semple, SJ, Dimsdale, JE, Ziegler, MG & Grant, I 2003, 'Emotional expressiveness, hostility and blood pressure in a longitudinal cohort of Alzheimer caregivers', *Journal of Psychosomatic Research*, vol. 54, no. 4, pp. 293-302.

Shorr, RI, Ray, WA, Daugherty, JR & Griffin, MR 1997, 'Antihypertensives and the risk of serious hypoglycemia in older persons using insulin or sulfonylureas', *Journal of the American Medical Association*, vol. 278, no. 1, pp. 40-43.

Sierksma, A, Lebrun, CE, van der Schouw, YT, Grobbee, DE, Lamberts, SW, Hendriks, HF & Bots, ML 2004, 'Alcohol consumption in relation to aortic stiffness and aortic wave reflections: a cross-sectional study in healthy postmenopausal women', *Arteriosclerosis, Thrombosis & Vascular Biology*, vol. 24, no. 2, pp. 342-348.

Siev-Ner, I, Gamus, D, Lerner-Geva, L & Achiron, A 2003, 'Reflexology treatment relieves symptoms of multiple sclerosis: a randomized controlled study', *Multiple Sclerosis*, vol. 9, no. 4, pp. 356-361.

Simons, LA, Simons, J, McCallum, J & Friedlander, Y 2005, 'Impact of smoking, diabetes and hypertension on survival time in the elderly: the Dubbo study', *Medical Journal of Australia*, vol. 182, pp. 219-222.

Singh, RB, Sharma, JP, Rastogi, V, Niaz, Ma & Singh, NK 1997, 'Prevalence and determinants of hypertension in the Indian social class and heart survey', *Journal of Human Hypertension*, vol. 11, no. 1, pp. 51-56.

Siscovick, DS, Raghunathan, TE, Psaty, BM, Koepsell, TD, Wicklund, KG, Lin, X, Cobb, L, Rautaharju, PM, Copass, MK & Wagner, EH 1994, 'Diuretic therapy for hypertension and the risk of primary cardiac arrest', *New England Journal of Medicine*, vol. 330, no. 26, pp. 1852-1857.

Skoumas, J, Pitsavos, C, Panagiotakos, DB, Chrysohoou, C, Zeimbekis, A, Papaioannou, I, Toutouza, M, Toutouzas, P & Stefanadis, C 2003, 'Physical activity, high density lipoprotein cholesterol and other lipids levels, in men and women from the ATTICA study', *Lipids in Health & Disease*, vol. 2, p. 3.

Smith, C, Birnbaum, G, Carter, JL, Greenstein, J & Lublin, FD 1994, 'Tizanidine treatment of spasticity caused by multiple sclerosis: results of a double-blind, placebo-controlled trial: US Tizanidine Study Group', *Neurology*, vol. 44, no. 11 Supp 19, pp. S34-42.

Smith-Barbaro, PA & Pucak, GJ 1983, 'Micronutrients and blood pressure: dietary fat and blood pressure', in *Annals of internal medicine: nutrition and blood pressure*

control, current status of dietary factors and hypertension, eds TA Kotchen & DA McCarron, The American College of Physicians, Philadelphia.

Spielberger, CD, Gorsuch, RL, Lushene, R, Vagg, PR & Jacobs, GA 1983, *The state-trait anxiety inventory*, Consulting psychologists press, California.

SPSS 12.0 for Windows 2003, CD-ROM, SPSS Inc., Chicago, Illinois.

Stefanick, ML, Mackey, S, Sheehan, M, Ellsworth, N, Haskell, WL & Wood, PD 1998, 'Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol', *New England Journal of Medicine*, vol. 339, no. 1, pp. 12-20.

Stephenson, N, Dalton, JA & Carlson, J 2003, 'The effect of foot reflexology on pain in patients with metastatic cancer', *Applied Nursing Research*, vol. 16, no. 4, pp. 284-286.

Stephenson, NLN, Weinrich, SP & Tavakoli, AS 2000, 'The effects of foot reflexology on anxiety and pain in patients with breast and lung cancer', *Oncology Nursing Forum*, vol. 27, no. 1, pp. 67-72.

Stephoe, A & Marmot, M 2005, 'Impaired cardiovascular recovery following stress predicts 3-year increases in blood pressure', *Journal of Hypertension*, vol. 23, no. 3, pp. 529-36.

Stevenson, CJ 1994, 'The psychophysiological effects of aromatherapy massage following cardiac surgery', *Complementary Therapies in Medicine*, vol. 2, pp. 27-35.

Stranges, S, Wu, T, Dorn, JM, Freudenheim, JL, Muti, P, Farinaro, E, Russell, M, Nochajski, TH & Trevisan, M 2004, 'Relationship of alcohol drinking pattern to risk of hypertension: a population-based study', *Hypertension*, vol. 44, no. 6, pp. 813-819.

Studer, M, Briel, M, Leimenstoll, B, Glass, TR & Bucher, HC 2005, 'Effect of different antilipidemic agents and diets on mortality', *Archives of Internal Medicine*, vol. 165, no. 7, pp. 725-730.

Subjaruean, P, Deeviset, K & Toieum, Y 2002, *Foot massage for health*, Samjarueanpanit, Bangkok, Thailand.

Swift, PA, Markandu, ND, Sagnella, GA, He, FJ & MacGregor, GA 2005, 'Modest salt reduction reduces blood pressure and urine protein excretion in black hypertensives: a randomized control trial', *Hypertension*, vol. 46, no. 2, pp. 308-312.

Swinscow, TDV 1997, *Statistics at square one*, viewed 16 March 2004, <<http://bmj.bmjournals.com/collections/statsbk/index.shtml>>.

Taomountain 2004, *An introduction to traditional Thai ayurvedic medicine*, viewed 3 August 2005, <<http://www.taomountain.org/thai-herbs/thai-ayurveda.html>>.

Tedesco, MA, Ratti, G, Di Salvo, G & Natale, F 2002, 'Does the angiotensin II receptor antagonist losartan improve cognitive function', *Drugs and Aging*, vol. 19, no. 10, pp. 723-732.

Testa, MA, Hollenberg, NK, Anderson, RB & Williams, GH 1991, 'Assessment of quality of life by patient and spouse during antihypertensive therapy with atenolol and nifedipine gastrointestinal therapeutic system', *American Journal of Hypertension*, vol. 4, pp. 363-373.

The American Heritage Stedman's Medical Dictionary 2002, *Valsalva maneuver*, viewed 2 November 2005, <<http://www.answers.com/topic/valsalva-maneuver>>.

The WHOQOL Group 1996, 'People and health: what quality of life?', *World Health Forum*, vol. 17, pp. 354-356.

The WHOQOL Group 1998, *RCMAR measurement tools*, viewed 3 December 2005, <<http://www.musc.edu/dfm/RCMAR/WHOQOL.html>>.

Theorell, T, de Faire, U, Johnson, J, Hall, E, Perski, A & Stewart, W 1991, 'Job strain and ambulatory blood pressure profiles', *Scandinavian Journal of Work, Environment and Health*, vol. 17, no. 6, pp. 380-385.

Thompson, GR 1990, *A handbook of hyperlipidaemia*. Current Science Ltd, London.

Timio, M, Saronio, P, Venanzi, S, Gentili, S, Verdura, C & Timio, F 1999, 'Blood pressure in nuns in a secluded order: a 30-year follow-up', *Mineral and Electrolyte Metabolism*, vol. 25, pp. 73-79.

Torrance, GW, Furlong, W, Feeny, D & Boyle, M 1995, 'Mult-attribute preference functions: health utilities index', *Pharmacoeconomics*, vol. 7, pp. 503-520.

Tovey, P 2002, 'A single-blind trial of reflexology for irritable bowel syndrome', *British Journal of General Practice*, vol. 52, pp. 19-23.

van Haselen, RA, Reiber, U, Nickel, I, Jakob, A & Fisher, PA 2004, 'Providing complementary and alternative medicine in primary care: the primary care workers' perspective', *Complementary Therapies in Medicine*, vol. 12, no. 1, pp. 6-16.

Vanitbanha, K 2003, *Using SPSS for Windows for data analysis*, 6th edn, Thammasan, Bangkok, Thailand.

Velasco, M, Lezama, E, Hernandez-Hernandez, R, Armas-Hernandez, MJ, Armas-Padilla, MC & Alcocer, L 2001, 'Semipersonalized psychological evaluation of quality of life in hypertensive patients during a trial with two calcium antagonists: a multicenter Latin-American Study (LASTLHY)', *International Journal of Clinical Pharmacology & Therapeutics*, vol. 39, no. 8, pp. 325-330.

Walker, C, Peterson, CL, Millen, N & Martin, C 2003, *Chronic illness: new perspectives and new directions*, Tertiary Press, Victoria.

Walker, MJ & Walker, JD 2003, *Healing massage: a simple approach*, Delmar Learning, New York.

Ware, JE & Gandek, B 1998, 'Overview of the SF-36 health survey and the international quality of life assessment (IQOLA) project', *Journal of Clinical Epidemiology*, vol. 51, no. 11, pp. 903-912.

Ware, JE & Kosinski, M 2001, 'Interpreting SF-36 summary health measures: a response', *Quality of Life Research*, vol. 10, pp. 405-413.

Ware, JE, Kosinski, M & Keller, SD 1994, *Physical and mental health summary scales: a user's manual*, The health institute, Boston, Massachusetts.

Ware, JE & Sherbourne, CD 1992, 'The MOS 36-item short-form health survey (SF-36) I: conceptual framework and item selection', *Medical Care*, vol. 30, pp. 473-483.

Wat Phra Baht Nam Phu n.d., *The Buddhist temple that cares for full-blown AIDS patients*, viewed 7 August 2005, <<http://www.aidstemple.th.org/index.html>>.

Welty, FK, Stuart, E, O'Meara, M & Huddleston, J 2002, 'Effect of addition of exercise to therapeutic lifestyle changes diet in enabling women and men with coronary heart disease to reach adult treatment panel III low-density lipoprotein cholesterol goal without lowering high-density lipoprotein cholesterol', *American Journal of Cardiology*, vol. 89, no. 10, pp. 1201-1204.

Whelton, SP, Chin, A, Xin, X & He, J 2002, 'Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials', *Annals of Internal Medicine*, vol. 136, no. 7, pp. 493-503.

White, LJ, Ferguson, MA, McCoy, SC & Kim, H 2003, 'Intramyocellular lipid changes in men and women during aerobic exercise: A¹ H-magnetic resonance spectroscopy study', *Journal of Clinical Endocrinology and Metabolism*, vol. 88, no. 12, pp. 5638-5643.

Whorwell, P 1984, 'Controlled trial of hypnotherapy in the treatment of severe refractory irritable bowel syndrome', *Lancet*, vol. 324, pp. 1232-1234.

Wiinberg, N, Hoegholm, A, Christensen, HR, Bang, LE, Mikkelsen, KL, Nielsen, PE, Svendsen, TL, Kampmann, JP, Madsen, NH & Bentzon, MW 1995, '24-h ambulatory blood pressure in 352 normal Danish subjects, related to age and gender', *American Journal of Hypertension*, vol. 8, no. 10 pt 1, pp. 978-986.

Wikstrand, J, Berglund, G & Tuomilehto, J 1991, 'Beta-blockade in the primary prevention of coronary heart disease in hypertensive patients: review of present evidence', *Circulation*, vol. 84, no. 6 Suppl, pp. VI93-VI100.

Williams, GH, Fisher, ND, Hunt, SC, Jeunemaitre, X, Hopkins, PN & Hollenberg, NK 2000, 'Genetics, pathophysiology, and pharmacological aspects: effects of gender and genotype on the phenotypic expression of nonmodulating essential hypertension', *Kidney International*, vol. 57, no. 4, pp. 1404-1407.

Williamson, J, White, A, Hart, A & Ernst, E 2002, 'Randomized controlled trial of reflexology for menopausal symptoms', *An International Journal of Obstetrics and Gynaecology*, vol. 109, no. 9, pp. 1050-1055.

Wolfgang, L 1984, *Psychological Perspectives of Essential Hypertension: Etiology, Maintenance, and Treatment*, Gasser & Cie AG, Switzerland.

Wongrattana, C 2001, *Tehcnique of using statistics for research*, 8th edn, Chulalongorn University, Bangkok, Thailand.

World Health Organization 1995, 'Economics of hypertension control', *Bulletin of the World Health Organization*, vol. 73, no. 4, pp. 417-424.

World Health Organization 1995, *WHOQOL-100: the 100 questions with response scales, field trial February 1995*, World Health Organization, Geneva.

World Health Organization 1996, *WHOQOL-BREF: introduction, administration, scoring and generic version of the assessment, field trial version December 1996*, World Health Organization, Geneva.

Wright, JM 2000, 'Choosing a first-line drug in the management of elevated blood pressure: what is the evidence? 2: β -blockers', *Canadian Medical Association Journal*, vol. 163, pp. 188-192.

Xin, X, He, J, Frontini, MG, Ogden, LG, Motsamai, OI & Whelton, PK 2001, 'Effects of alcohol reduction on blood pressure: a meta-analysis of randomized controlled trials', *Hypertension*, vol. 38, no. 5, pp. 1112-1117.

Yang, JH 2005, 'The effects of foot reflexology on nausea, vomiting and fatigue of breast cancer patients undergoing chemotherapy', *Daehan Ganho Haghoeji*, vol. 35, no. 1, pp. 177-185.

Yu, HC, Burrell, LM, Black, MJ, Wu, LL, Dilley, RJ, Cooper, ME & Johnston, CI 1998, 'Salt induces myocardial and renal fibrosis in normotensive and hypertensive rats', *Circulation*, vol. 98, no. 23, pp. 2621-2628.

Zanchetti, A, Omboni, S, La Commare, P, De Cesaris, R & Palatini, P 2001, 'Efficacy, tolerability, and impact on quality of life of long-term treatment with manidipine or amlodipine in patients with essential hypertension', *Journal of Cardiovascular Pharmacology*, vol. 38, no. 4, pp. 642-650.

Zilkens, RR, Burke, V, Hodgson, JM, Barden, A, Beilin, LJ & Puddey, IB 2005, 'Red wine and beer elevate blood pressure in normotensive men', *Hypertension*, vol. 45, no. 5, pp. 874-879.