

Reference

- Acheson, A., Conover, J. C., Fandl, J. P., DeChiara, T. M., Russell, M., Thadani, A., Squinto, S. P., Yancopoulos, G. D. and Lindsay, R. M. (1995) A BDNF autocrine loop in adult sensory neurons prevents cell death. *Nature*, **374**, 450-453.
- Agthong, S., Kaewsema, A., Tanomsridejchai, N. and Chentanez, V. (2006) Activation of MAPK ERK in peripheral nerve after injury. *BMC Neurosci*, **7**, 45.
- Aloyz, R. S., Bamji, S. X., Pozniak, C. D., Toma, J. G., Atwal, J., Kaplan, D. R. and Miller, F. D. (1998) p53 is essential for developmental neuron death as regulated by the TrkA and p75 neurotrophin receptors. *J Cell Biol*, **143**, 1691-1703.
- Altar, C. A., Cai, N., Bliven, T., Juhasz, M., Conner, J. M., Acheson, A. L., Lindsay, R. M. and Wiegand, S. J. (1997) Anterograde transport of brain-derived neurotrophic factor and its role in the brain. *Nature*, **389**, 856-860.
- Altar, C. A. and DiStefano, P. S. (1998) Neurotrophin trafficking by anterograde transport. *Trends Neurosci*, **21**, 433-437.
- Baas, P. W., Black, M. M. and Bunker, G. A. (1989) Changes in microtubule polarity orientation during the development of hippocampal neurons in culture. *J Cell Biol*, **109**, 3085-3094.
- Balkowiec, A. and Katz, D. M. (2002) Cellular mechanisms regulating activity-dependent release of native brain-derived neurotrophic factor from

- hippocampal neurons. *J Neurosci*, **22**, 10399-10407.
- Bamji, S. X., Majdan, M., Pozniak, C. D., Belliveau, D. J., Aloyz, R., Kohn, J., Causing, C. G. and Miller, F. D. (1998) The p75 neurotrophin receptor mediates neuronal apoptosis and is essential for naturally occurring sympathetic neuron death. *J Cell Biol*, **140**, 911-923.
- Baquet, Z. C., Gorski, J. A. and Jones, K. R. (2004) Early striatal dendrite deficits followed by neuron loss with advanced age in the absence of anterograde cortical brain-derived neurotrophic factor. *J Neurosci*, **24**, 4250-4258.
- Barde, Y. A. (1990) The nerve growth factor family. *Prog Growth Factor Res*, **2**, 237-248.
- Bath, K. G. and Lee, F. S. (2006) Variant BDNF (Val66Met) impact on brain structure and function. *Cogn Affect Behav Neurosci*, **6**, 79-85.
- Beattie, M. S., Harrington, A. W., Lee, R., Kim, J. Y., Boyce, S. L., Longo, F. M., Bresnahan, J. C., Hempstead, B. L. and Yoon, S. O. (2002) ProNGF induces p75-mediated death of oligodendrocytes following spinal cord injury. *Neuron*, **36**, 375-386.
- Berkemeier, L. R., Winslow, J. W., Kaplan, D. R., Nikolics, K., Goeddel, D. V. and Rosenthal, A. (1991) Neurotrophin-5: a novel neurotrophic factor that activates trk and trkB. *Neuron*, **7**, 857-866.
- Berridge, M. J. (1998) Neuronal calcium signaling. *Neuron*, **21**, 13-26.

- Bertaux, F., Sharp, A. H., Ross, C. A., Lehrach, H., Bates, G. P. and Wanker, E. (1998) HAP1-huntingtin interactions do not contribute to the molecular pathology in Huntington's disease transgenic mice. *FEBS Lett.*, **426**, 229-232.
- Bhattacharyya, A., Watson, F. L., Pomeroy, S. L., Zhang, Y. Z., Stiles, C. D. and Segal, R. A. (2002) High-resolution imaging demonstrates dynein-based vesicular transport of activated Trk receptors. *J Neurobiol.*, **51**, 302-312.
- Bibel, M., Hoppe, E. and Barde, Y. A. (1999) Biochemical and functional interactions between the neurotrophin receptors trk and p75NTR. *Embo J.*, **18**, 616-622.
- Bilderback, T. R., Gazula, V. R., Lisanti, M. P. and Dobrowsky, R. T. (1999) Caveolin interacts with Trk A and p75(NTR) and regulates neurotrophin signaling pathways. *J Biol Chem.*, **274**, 257-263.
- Binder, D. K., Croll, S. D., Gall, C. M. and Scharfman, H. E. (2001) BDNF and epilepsy: too much of a good thing? *Trends Neurosci.*, **24**, 47-53.
- Binder, D. K. and Scharfman, H. E. (2004) Brain-derived neurotrophic factor. *Growth Factors*, **22**, 123-131.
- Blochl, A. and Thoenen, H. (1995) Characterization of nerve growth factor (NGF) release from hippocampal neurons: evidence for a constitutive and an unconventional sodium-dependent regulated pathway. *Eur J Neurosci.*, **7**, 1220-1228.
- Block-Galarza, J., Chase, K. O., Sapp, E., Vaughn, K. T., Vallee, R. B.,

- DiFiglia, M. and Aronin, N. (1997) Fast transport and retrograde movement of huntingtin and HAP 1 in axons. *Neuroreport*, **8**, 2247-2251.
- Bonifacino, J. S., Marks, M. S., Ohno, H. and Kirchhausen, T. (1996) Mechanisms of signal-mediated protein sorting in the endocytic and secretory pathways. *Proc Assoc Am Physicians*, **108**, 285-295.
- Bossy-Wetzel, E., Schwarzenbacher, R. and Lipton, S. A. (2004) Molecular pathways to neurodegeneration. *Nat Med*, **10 Suppl**, S2-9.
- Bothwell, M. (1995) Functional interactions of neurotrophins and neurotrophin receptors. *Annu Rev Neurosci*, **18**, 223-253.
- Boukhaddaoui, H., Sieso, V., Scamps, F. and Valmier, J. (2001) An activity-dependent neurotrophin-3 autocrine loop regulates the phenotype of developing hippocampal pyramidal neurons before target contact. *J Neurosci*, **21**, 8789-8797.
- Bradshaw, R. A., Murray-Rust, J., Ibanez, C. F., McDonald, N. Q., Lapatto, R. and Blundell, T. L. (1994) Nerve growth factor: structure/function relationships. *Protein Sci*, **3**, 1901-1913.
- Brady, R., Zaidi, S. I., Mayer, C. and Katz, D. M. (1999) BDNF is a target-derived survival factor for arterial baroreceptor and chemoafferent primary sensory neurons. *J Neurosci*, **19**, 2131-2142.
- Bronfman, F. C., Tcherpakov, M., Jovin, T. M. and Fainzilber, M. (2003) Ligand-induced internalization of the p75 neurotrophin receptor: a slow route

- to the signaling endosome. *J Neurosci*, **23**, 3209-3220.
- Brunet, A., Datta, S. R. and Greenberg, M. E. (2001) Transcription-dependent and -independent control of neuronal survival by the PI3K-Akt signaling pathway. *Curr Opin Neurobiol*, **11**, 297-305.
- Burgess, S. A., Walker, M. L., Sakakibara, H., Knight, P. J. and Oiwa, K. (2003) Dynein structure and power stroke. *Nature*, **421**, 715-718.
- Burnett, M. G. and Zager, E. L. (2004) Pathophysiology of peripheral nerve injury: a brief review. *Neurosurg Focus*, **16**, E1.
- Butowt, R. and von Bartheld, C. S. (2001) Sorting of internalized neurotrophins into an endocytic transcytosis pathway via the Golgi system: Ultrastructural analysis in retinal ganglion cells. *J Neurosci*, **21**, 8915-8930.
- Buttigieg, H., Kawaja, M. D. and Fahnstock, M. (2007) Neurotrophic activity of proNGF in vivo. *Exp Neurol*, **204**, 832-835.
- Campenot, R. B. and MacInnis, B. L. (2004) Retrograde transport of neurotrophins: fact and function. *J Neurobiol*, **58**, 217-229.
- Carter, B. D., Dechant, G., Frade, J. M., Kaltschmidt, C. and Barde, Y. A. (1996) Neurotrophins and their p75 receptor. *Cold Spring Harb Symp Quant Biol*, **61**, 407-415.
- Cattaneo, E., Zuccato, C. and Tartari, M. (2005) Normal huntingtin function: an alternative approach to Huntington's disease. *Nat Rev Neurosci*, **6**, 919-930.
- Chan, E. Y., Nasir, J., Gutekunst, C. A. et al. (2002) Targeted disruption of

- Huntingtin-associated protein-1 (Hap1) results in postnatal death due to depressed feeding behavior. *Hum Mol Genet*, **11**, 945-959.
- Chang, M. S., Arevalo, J. C. and Chao, M. V. (2004) Ternary complex with Trk, p75, and an ankyrin-rich membrane spanning protein. *J Neurosci Res*, **78**, 186-192.
- Chao, M. V. and Bothwell, M. (2002) Neurotrophins: to cleave or not to cleave. *Neuron*, **33**, 9-12.
- Charrin, B. C., Saudou, F. and Humbert, S. (2005) Axonal transport failure in neurodegenerative disorders: the case of Huntington's disease. *Pathol Biol (Paris)*, **53**, 189-192.
- Chen, Y. W. (2003) Local protein unfolding and pathogenesis of polyglutamine-expansion diseases. *Proteins*, **51**, 68-73.
- Chen, Z. Y., Ieraci, A., Teng, H., Dall, H., Meng, C. X., Herrera, D. G., Nykjaer, A., Hempstead, B. L. and Lee, F. S. (2005) Sortilin controls intracellular sorting of brain-derived neurotrophic factor to the regulated secretory pathway. *J Neurosci*, **25**, 6156-6166.
- Chen, Z. Y., Patel, P. D., Sant, G., Meng, C. X., Teng, K. K., Hempstead, B. L. and Lee, F. S. (2004) Variant brain-derived neurotrophic factor (BDNF) (Met66) alters the intracellular trafficking and activity-dependent secretion of wild-type BDNF in neurosecretory cells and cortical neurons. *J Neurosci*, **24**, 4401-4411.

- Choi, N. S., Yoo, K. H., Yoon, K. S., Maeng, P. J. and Kim, S. H. (2004a) Nano-scale proteomics approach using two-dimensional fibrin zymography combined with fluorescent SYPRO ruby dye. *J Biochem Mol Biol*, **37**, 298-303.
- Choi, S. Y., Hwang, J. J. and Koh, J. Y. (2004b) NR2A induction and NMDA receptor-dependent neuronal death by neurotrophin-4/5 in cortical cell culture. *J Neurochem*, **88**, 708-716.
- Cohen, S. and Levi-Montalcini, R. (1957) Purification and properties of a nerve growth-promoting factor isolated from mouse sarcoma 180. *Cancer Res*, **17**, 15-20.
- Colomer, V., Engelender, S., Sharp, A. H., Duan, K., Cooper, J. K., Lanahan, A., Lyford, G., Worley, P. and Ross, C. A. (1997) Huntington-associated protein 1 (HAP1) binds to a Trio-like polypeptide, with a rac1 guanine nucleotide exchange factor domain. *Hum Mol Genet*, **6**, 1519-1525.
- Conner, J. M., Lauterborn, J. C., Yan, Q., Gall, C. M. and Varon, S. (1997) Distribution of brain-derived neurotrophic factor (BDNF) protein and mRNA in the normal adult rat CNS: evidence for anterograde axonal transport. *J Neurosci*, **17**, 2295-2313.
- Coppola, V., Kucera, J., Palko, M. E., Martinez-De Velasco, J., Lyons, W. E., Fritzsch, B. and Tessarollo, L. (2001) Dissection of NT3 functions in vivo by gene replacement strategy. *Development*, **128**, 4315-4327.

Reference

- Corbit, K. C., Foster, D. A. and Rosner, M. R. (1999) Protein kinase C δ mediates neurogenic but not mitogenic activation of mitogen-activated protein kinase in neuronal cells. *Mol Cell Biol*, **19**, 4209-4218.
- Crowley, C., Spencer, S. D., Nishimura, M. C. et al. (1994) Mice lacking nerve growth factor display perinatal loss of sensory and sympathetic neurons yet develop basal forebrain cholinergic neurons. *Cell*, **76**, 1001-1011.
- Davies, S. and Ramsden, D. B. (2001) Huntington's disease. *Mol Pathol*, **54**, 409-413.
- Deacon, S. W., Serpinskaya, A. S., Vaughan, P. S., Lopez Fanarraga, M., Vernos, I., Vaughan, K. T. and Gelfand, V. I. (2003) Dynactin is required for bidirectional organelle transport. *J Cell Biol*, **160**, 297-301.
- Dechant, G. (2001) Molecular interactions between neurotrophin receptors. *Cell Tissue Res*, **305**, 229-238.
- Dechant, G. and Barde, Y. A. (2002) The neurotrophin receptor p75(NTR): novel functions and implications for diseases of the nervous system. *Nat Neurosci*, **5**, 1131-1136.
- Dechant, G. and Neumann, H. (2002) Neurotrophins. *Adv Exp Med Biol*, **513**, 303-334.
- Dechant, G., Tsoulfas, P., Parada, L. F. and Barde, Y. A. (1997) The neurotrophin receptor p75 binds neurotrophin-3 on sympathetic neurons with high affinity and specificity. *J Neurosci*, **17**, 5281-5287.

- del Toro, D., Canals, J. M., Gines, S., Kojima, M., Egea, G. and Alberch, J. (2006) Mutant huntingtin impairs the post-Golgi trafficking of brain-derived neurotrophic factor but not its Val66Met polymorphism. *J Neurosci*, **26**, 12748-12757.
- Dell, K. R. (2003) Dynactin polices two-way organelle traffic. *J Cell Biol*, **160**, 291-293.
- DiFiglia, M., Sapp, E., Chase, K. et al. (1995) Huntingtin is a cytoplasmic protein associated with vesicles in human and rat brain neurons. *Neuron*, **14**, 1075-1081.
- Domeniconi, M., Hempstead, B. L. and Chao, M. V. (2007) Pro-NGF secreted by astrocytes promotes motor neuron cell death. *Mol Cell Neurosci*, **34**, 271-279.
- Dragatsis, I., Zeitlin, S. and Dietrich, P. (2004) Huntingtin-associated protein 1 (Hap1) mutant mice bypassing the early postnatal lethality are neuroanatomically normal and fertile but display growth retardation. *Hum Mol Genet*, **13**, 3115-3125.
- Du, Y., Fischer, T. Z., Clinton-Luke, P., Lercher, L. D. and Dreyfus, C. F. (2006) Distinct effects of p75 in mediating actions of neurotrophins on basal forebrain oligodendrocytes. *Mol Cell Neurosci*, **31**, 366-375.
- Edwards, R. H., Selby, M. J., Garcia, P. D. and Rutter, W. J. (1988) Processing of the native nerve growth factor precursor to form biologically active nerve

- growth factor. *J Biol Chem*, **263**, 6810-6815.
- Egan, M. F., Kojima, M., Callicott, J. H. et al. (2003) The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell*, **112**, 257-269.
- ElShamy, W. M. and Ernfors, P. (1997) Brain-derived neurotrophic factor, neurotrophin-3, and neurotrophin-4 complement and cooperate with each other sequentially during visceral neuron development. *J Neurosci*, **17**, 8667-8675.
- Engelender, S., Sharp, A. H., Colomer, V., Tokito, M. K., Lanahan, A., Worley, P., Holzbaur, E. L. and Ross, C. A. (1997) Huntington-associated protein 1 (HAP1) interacts with the p150Glued subunit of dynactin. *Hum Mol Genet*, **6**, 2205-2212.
- Erickson, J. T., Conover, J. C., Borday, V., Champagnat, J., Barbacid, M., Yancopoulos, G. and Katz, D. M. (1996) Mice lacking brain-derived neurotrophic factor exhibit visceral sensory neuron losses distinct from mice lacking NT4 and display a severe developmental deficit in control of breathing. *J Neurosci*, **16**, 5361-5371.
- Erickson, J. W. and Cerione, R. A. (2004) Structural elements, mechanism, and evolutionary convergence of Rho protein-guanine nucleotide exchange factor complexes. *Biochemistry*, **43**, 837-842.
- Ernfors, P., Ibanez, C. F., Ebendal, T., Olson, L. and Persson, H. (1990) Molecular cloning and neurotrophic activities of a protein with structural

- similarities to nerve growth factor: developmental and topographical expression in the brain. *Proc Natl Acad Sci U S A*, **87**, 5454-5458.
- Ernfors, P., Lee, K. F. and Jaenisch, R. (1994) Mice lacking brain-derived neurotrophic factor develop with sensory deficits. *Nature*, **368**, 147-150.
- Fahnstock, M., Michalski, B., Xu, B. and Coughlin, M. D. (2001) The precursor pro-nerve growth factor is the predominant form of nerve growth factor in brain and is increased in Alzheimer's disease. *Mol Cell Neurosci*, **18**, 210-220.
- Fahnstock, M., Yu, G., Michalski, B., Mathew, S., Colquhoun, A., Ross, G. M. and Coughlin, M. D. (2004) The nerve growth factor precursor proNGF exhibits neurotrophic activity but is less active than mature nerve growth factor. *J Neurochem*, **89**, 581-592.
- Farhadi, H. F., Mowla, S. J., Petrecca, K., Morris, S. J., Seidah, N. G. and Murphy, R. A. (2000) Neurotrophin-3 sorts to the constitutive secretory pathway of hippocampal neurons and is diverted to the regulated secretory pathway by coexpression with brain-derived neurotrophic factor. *J Neurosci*, **20**, 4059-4068.
- Farinas, I. (1999) Neurotrophin actions during the development of the peripheral nervous system. *Microsc Res Tech*, **45**, 233-242.
- Farinas, I., Wilkinson, G. A., Backus, C., Reichardt, L. F. and Patapoutian, A. (1998) Characterization of neurotrophin and Trk receptor functions in

- developing sensory ganglia: direct NT-3 activation of TrkB neurons in vivo. *Neuron*, **21**, 325-334.
- Farinas, I., Yoshida, C. K., Backus, C. and Reichardt, L. F. (1996) Lack of neurotrophin-3 results in death of spinal sensory neurons and premature differentiation of their precursors. *Neuron*, **17**, 1065-1078.
- Fayard, B., Loeffler, S., Weis, J., Vogelin, E. and Kruttgen, A. (2005) The secreted brain-derived neurotrophic factor precursor pro-BDNF binds to TrkB and p75NTR but not to TrkA or TrkC. *J Neurosci Res*, **80**, 18-28.
- Ferrer, I. (1999) [Neuronal death mechanisms in cerebral ischemia]. *Rev Neurol*, **29**, 515-521.
- Ferrer, I., Goutan, E., Marin, C., Rey, M. J. and Ribalta, T. (2000) Brain-derived neurotrophic factor in Huntington disease. *Brain Res*, **866**, 257-261.
- Foehr, E. D., Lin, X., O'Mahony, A., Geleziunas, R., Bradshaw, R. A. and Greene, W. C. (2000) NF-kappa B signaling promotes both cell survival and neurite process formation in nerve growth factor-stimulated PC12 cells. *J Neurosci*, **20**, 7556-7563.
- Frostick, S. P., Yin, Q. and Kemp, G. J. (1998) Schwann cells, neurotrophic factors, and peripheral nerve regeneration. *Microsurgery*, **18**, 397-405.
- Fryer, H. J., Wolf, D. H., Knox, R. J., Strittmatter, S. M., Pennica, D., O'Leary, R. M., Russell, D. S. and Kalb, R. G. (2000) Brain-derived neurotrophic factor induces excitotoxic sensitivity in cultured embryonic rat spinal motor neurons

- through activation of the phosphatidylinositol 3-kinase pathway. *J Neurochem*, **74**, 582-595.
- Gabriely, G., Kama, R. and Gerst, J. E. (2007) Involvement of specific COPI subunits in protein sorting from the late endosome to the vacuole in yeast. *Mol Cell Biol*, **27**, 526-540.
- Gaglio, T., Dionne, M. A. and Compton, D. A. (1997) Mitotic spindle poles are organized by structural and motor proteins in addition to centrosomes. *J Cell Biol*, **138**, 1055-1066.
- Garner, C. C., Nash, J. and Huganir, R. L. (2000) PDZ domains in synapse assembly and signalling. *Trends Cell Biol*, **10**, 274-280.
- Garzon, D. J. and Fahnestock, M. (2007) Oligomeric amyloid decreases basal levels of brain-derived neurotrophic factor (BDNF) mRNA via specific downregulation of BDNF transcripts IV and V in differentiated human neuroblastoma cells. *J Neurosci*, **27**, 2628-2635.
- Gatzinsky, K. P., Haugland, R. P., Thrasivoulou, C., Orike, N., Budi-Santoso, A. W. and Cowen, T. (2001) p75 and TrkA receptors are both required for uptake of NGF in adult sympathetic neurons: use of a novel fluorescent NGF conjugate. *Brain Res*, **920**, 226-238.
- Gauthier, L. R., Charrin, B. C., Borrell-Pages, M. et al. (2004) Huntington controls neurotrophic support and survival of neurons by enhancing BDNF vesicular transport along microtubules. *Cell*, **118**, 127-138.

- Gines, S., Bosch, M., Marco, S., Gavalda, N., Diaz-Hernandez, M., Lucas, J. J., Canals, J. M. and Alberch, J. (2006) Reduced expression of the TrkB receptor in Huntington's disease mouse models and in human brain. *Eur J Neurosci*, **23**, 649-658.
- Ginty, D. D. and Segal, R. A. (2002) Retrograde neurotrophin signaling: Trk-ing along the axon. *Curr Opin Neurobiol*, **12**, 268-274.
- Goodman, L. J., Valverde, J., Lim, F., Geschwind, M. D., Federoff, H. J., Geller, A. I. and Hefti, F. (1996) Regulated release and polarized localization of brain-derived neurotrophic factor in hippocampal neurons. *Mol Cell Neurosci*, **7**, 222-238.
- Goodman, S. R., Zimmer, W. E., Clark, M. B., Zagon, I. S., Barker, J. E. and Bloom, M. L. (1995) Brain spectrin: of mice and men. *Brain Res Bull*, **36**, 593-606.
- Gray, E. G. and Whittaker, V. P. (1962) The isolation of nerve endings from brain: an electron-microscopic study of cell fragments derived by homogenization and centrifugation. *J Anat*, **96**, 79-88.
- Griesbeck, O., Canossa, M., Campana, G., Gartner, A., Hoener, M. C., Nawa, H., Kolbeck, R. and Thoenen, H. (1999) Are there differences between the secretion characteristics of NGF and BDNF? Implications for the modulatory role of neurotrophins in activity-dependent neuronal plasticity. *Microsc Res Tech*, **45**, 262-275.

- Gross, S. P., Welte, M. A., Block, S. M. and Wieschaus, E. F. (2002) Coordination of opposite-polarity microtubule motors. *J Cell Biol*, **156**, 715-724.
- Gu, F., Crump, C. M. and Thomas, G. (2001) Trans-Golgi network sorting. *Cell Mol Life Sci*, **58**, 1067-1084.
- Gueorguiev, V. D., Zeman, R. J., Hiremagalur, B., Menezes, A. and Sabban, E. L. (1999) Differing temporal roles of Ca²⁺ and cAMP in nicotine-elicited elevation of tyrosine hydroxylase mRNA. *Am J Physiol*, **276**, C54-65.
- Guillaud, L., Setou, M. and Hirokawa, N. (2003) KIF17 dynamics and regulation of NR2B trafficking in hippocampal neurons. *J Neurosci*, **23**, 131-140.
- Gunawardena, S. and Goldstein, L. S. (2004a) Cargo-carrying motor vehicles on the neuronal highway: transport pathways and neurodegenerative disease. In: *J Neurobiol*, Vol. 58, pp. 258-271.
- Gunawardena, S. and Goldstein, L. S. (2004b) Cargo-carrying motor vehicles on the neuronal highway: transport pathways and neurodegenerative disease. *J Neurobiol*, **58**, 258-271.
- Gutekunst, C. A., Li, S. H., Yi, H., Ferrante, R. J., Li, X. J. and Hersch, S. M. (1998) The cellular and subcellular localization of huntingtin-associated protein 1 (HAP1): comparison with huntingtin in rat and human. *J Neurosci*, **18**, 7674-7686.

- Hamanoue, M., Middleton, G., Wyatt, S., Jaffray, E., Hay, R. T. and Davies, A. M. (1999) p75-mediated NF-kappaB activation enhances the survival response of developing sensory neurons to nerve growth factor. *Mol Cell Neurosci*, **14**, 28-40.
- Hariri, A. R., Goldberg, T. E., Mattay, V. S., Kolachana, B. S., Callicott, J. H., Egan, M. F. and Weinberger, D. R. (2003) Brain-derived neurotrophic factor val66met polymorphism affects human memory-related hippocampal activity and predicts memory performance. *J Neurosci*, **23**, 6690-6694.
- Harrington, A. W., Leiner, B., Blechschmitt, C. et al. (2004) Secreted proNGF is a pathophysiological death-inducing ligand after adult CNS injury. *Proc Natl Acad Sci U S A*, **101**, 6226-6230.
- Hashimoto, K. (2007) BDNF variant linked to anxiety-related behaviors. *Bioessays*, **29**, 116-119.
- Hempstead, B. L. (2006) Dissecting the diverse actions of pro- and mature neurotrophins. *Curr Alzheimer Res*, **3**, 19-24.
- Henderson, C. E., Camu, W., Mettling, C. et al. (1993) Neurotrophins promote motor neuron survival and are present in embryonic limb bud. *Nature*, **363**, 266-270.
- Hendry, I. A., Stach, R. and Herrup, K. (1974a) Characteristics of the retrograde axonal transport system for nerve growth factor in the sympathetic nervous system. *Brain Res*, **82**, 117-128.

- Hendry, I. A., Stockel, K., Thoenen, H. and Iversen, L. L. (1974b) The retrograde axonal transport of nerve growth factor. *Brain Res*, **68**, 103-121.
- Heymach, J. V., Jr. and Shooter, E. M. (1995) The biosynthesis of neurotrophin heterodimers by transfected mammalian cells. *J Biol Chem*, **270**, 12297-12304.
- Hibbert, A. P., Morris, S. J., Seidah, N. G. and Murphy, R. A. (2003) Neurotrophin-4, alone or heterodimerized with brain-derived neurotrophic factor, is sorted to the constitutive secretory pathway. *J Biol Chem*, **278**, 48129-48136.
- Hirokawa, N. and Takemura, R. (2004) Molecular motors in neuronal development, intracellular transport and diseases. *Curr Opin Neurobiol*, **14**, 564-573.
- Hirokawa, N. and Takemura, R. (2005) Molecular motors and mechanisms of directional transport in neurons. *Nat Rev Neurosci*, **6**, 201-214.
- Hohn, A., Leibrock, J., Bailey, K. and Barde, Y. A. (1990) Identification and characterization of a novel member of the nerve growth factor/brain-derived neurotrophic factor family. *Nature*, **344**, 339-341.
- Hongo, T., Hakuba, A., Shiota, K. and Naruse, I. (2000) Suckling dysfunction caused by defects in the olfactory system in genetic arhinencephaly mice. *Biol Neonate*, **78**, 293-299.
- Horch, H. W. and Katz, L. C. (2002) BDNF release from single cells elicits local dendritic growth in nearby neurons. *Nat Neurosci*, **5**, 1177-1184.

- Howells, D. W., Porritt, M. J., Wong, J. Y., Batchelor, P. E., Kalnins, R., Hughes, A. J. and Donnan, G. A. (2000) Reduced BDNF mRNA expression in the Parkinson's disease substantia nigra. *Exp Neurol*, **166**, 127-135.
- Huang, E. J. and Reichardt, L. F. (2001) Neurotrophins: roles in neuronal development and function. *Annu Rev Neurosci*, **24**, 677-736.
- Huang, E. J. and Reichardt, L. F. (2003) Trk receptors: roles in neuronal signal transduction. *Annu Rev Biochem*, **72**, 609-642.
- Humbert, S. and Saudou, F. (2005) [Huntington's disease: intracellular signaling pathways and neuronal death]. *J Soc Biol*, **199**, 247-251.
- Hung, A. Y. and Sheng, M. (2002) PDZ domains: structural modules for protein complex assembly. *J Biol Chem*, **277**, 5699-5702.
- Jo, K., Derin, R., Li, M. and Bredt, D. S. (1999) Characterization of MALS/Velis-1, -2, and -3: a family of mammalian LIN-7 homologs enriched at brain synapses in association with the postsynaptic density-95/NMDA receptor postsynaptic complex. *J Neurosci*, **19**, 4189-4199.
- Johnson, E. M., Jr., Taniuchi, M., Clark, H. B., Springer, J. E., Koh, S., Tayrien, M. W. and Loy, R. (1987) Demonstration of the retrograde transport of nerve growth factor receptor in the peripheral and central nervous system. *J Neurosci*, **7**, 923-929.
- Johnson, F., Hohmann, S. E., DiStefano, P. S. and Bottjer, S. W. (1997) Neurotrophins suppress apoptosis induced by deafferentation of an avian

- motor-cortical region. *J Neurosci*, **17**, 2101-2111.
- Jones, K. R., Farinas, I., Backus, C. and Reichardt, L. F. (1994) Targeted disruption of the BDNF gene perturbs brain and sensory neuron development but not motor neuron development. *Cell*, **76**, 989-999.
- Kalb, R. (2005) The protean actions of neurotrophins and their receptors on the life and death of neurons. *Trends Neurosci*, **28**, 5-11.
- Kanai, Y., Dohmae, N. and Hirokawa, N. (2004) Kinesin transports RNA: isolation and characterization of an RNA-transporting granule. *Neuron*, **43**, 513-525.
- Kanai, Y., Okada, Y., Tanaka, Y., Harada, A., Terada, S. and Hirokawa, N. (2000) KIF5C, a novel neuronal kinesin enriched in motor neurons. *J Neurosci*, **20**, 6374-6384.
- Kaplan, D. R. and Miller, F. D. (2000) Neurotrophin signal transduction in the nervous system. *Curr Opin Neurobiol*, **10**, 381-391.
- Karlin, S. and Burge, C. (1996) Trinucleotide repeats and long homopeptides in genes and proteins associated with nervous system disease and development. *Proc Natl Acad Sci U S A*, **93**, 1560-1565.
- Katoh-Semba, R., Asano, T., Ueda, H., Morishita, R., Takeuchi, I. K., Inaguma, Y. and Kato, K. (2002) Riluzole enhances expression of brain-derived neurotrophic factor with consequent proliferation of granule precursor cells in the rat hippocampus. *Faseb J*, **16**, 1328-1330.

Reference

- Katz, L. C. and Shatz, C. J. (1996) Synaptic activity and the construction of cortical circuits. *Science*, **274**, 1133-1138.
- Kells, A. P., Fong, D. M., Dragunow, M., During, M. J., Young, D. and Connor, B. (2004) AAV-mediated gene delivery of BDNF or GDNF is neuroprotective in a model of Huntington disease. *Mol Ther*, **9**, 682-688.
- Kim, M., Velier, J., Chase, K. et al. (1999) Forskolin and dopamine D1 receptor activation increase huntingtin's association with endosomes in immortalized neuronal cells of striatal origin. *Neuroscience*, **89**, 1159-1167.
- Kittler, J. T., Thomas, P., Tretter, V., Bogdanov, Y. D., Haucke, V., Smart, T. G. and Moss, S. J. (2004) Huntingtin-associated protein 1 regulates inhibitory synaptic transmission by modulating gamma-aminobutyric acid type A receptor membrane trafficking. *Proc Natl Acad Sci U S A*, **101**, 12736-12741.
- Klein, M., Hempstead, B. L. and Teng, K. K. (2005) Activation of STAT5-dependent transcription by the neurotrophin receptor Trk. *J Neurobiol*, **63**, 159-171.
- Klein, R., Smeyne, R. J., Wurst, W., Long, L. K., Auerbach, B. A., Joyner, A. L. and Barbacid, M. (1993) Targeted disruption of the trkB neurotrophin receptor gene results in nervous system lesions and neonatal death. *Cell*, **75**, 113-122.
- Kohara, K., Kitamura, A., Morishima, M. and Tsumoto, T. (2001) Activity-dependent transfer of brain-derived neurotrophic factor to postsynaptic

- neurons. *Science*, **291**, 2419-2423.
- Kojima, M., Takei, N., Numakawa, T., Ishikawa, Y., Suzuki, S., Matsumoto, T., Katoh-Semba, R., Nawa, H. and Hatanaka, H. (2001) Biological characterization and optical imaging of brain-derived neurotrophic factor-green fluorescent protein suggest an activity-dependent local release of brain-derived neurotrophic factor in neurites of cultured hippocampal neurons. *J Neurosci Res*, **64**, 1-10.
- Kolbeck, R., Jungbluth, S. and Barde, Y. A. (1994) Characterisation of neurotrophin dimers and monomers. *Eur J Biochem*, **225**, 995-1003.
- Komada, M. and Kitamura, N. (2001) Hrs and hbp: possible regulators of endocytosis and exocytosis. *Biochem Biophys Res Commun*, **281**, 1065-1069.
- Kong, H., Boulter, J., Weber, J. L., Lai, C. and Chao, M. V. (2001) An evolutionarily conserved transmembrane protein that is a novel downstream target of neurotrophin and ephrin receptors. *J Neurosci*, **21**, 176-185.
- Korschning, S. (1993) The neurotrophic factor concept: a reexamination. *J Neurosci*, **13**, 2739-2748.
- Lalli, G. and Schiavo, G. (2002) Analysis of retrograde transport in motor neurons reveals common endocytic carriers for tetanus toxin and neurotrophin receptor p75NTR. *J Cell Biol*, **156**, 233-239.
- Larson, M. A. and Stein, B. E. (1984) The use of tactile and olfactory cues in neonatal orientation and localization of the nipple. *Dev Psychobiol*, **17**, 423-

436.

Lee, P., Zhuo, H. and Helke, C. J. (2001a) Axotomy alters neurotrophin and neurotrophin receptor mRNAs in the vagus nerve and nodose ganglion of the rat. *Brain Res Mol Brain Res*, **87**, 31-41.

Lee, R., Kermani, P., Teng, K. K. and Hempstead, B. L. (2001b) Regulation of cell survival by secreted proneurotrophins. *Science*, **294**, 1945-1948.

LeMaster, A. M., Krimm, R. F., Davis, B. M., Noel, T., Forbes, M. E., Johnson, J. E. and Albers, K. M. (1999) Overexpression of brain-derived neurotrophic factor enhances sensory innervation and selectively increases neuron number. *J Neurosci*, **19**, 5919-5931.

Lesuisse, C. and Martin, L. J. (2002) Long-term culture of mouse cortical neurons as a model for neuronal development, aging, and death. *J Neurobiol*, **51**, 9-23.

Levi-Montalcini, R. (1949) The development to the acoustico-vestibular centers in the chick embryo in the absence of the afferent root fibers and of descending fiber tracts. *J Comp Neurol*, **91**, 209-241, illust, incl 203 pl.

Levi-Montalcini, R., Skaper, S. D., Dal Toso, R., Petrelli, L. and Leon, A. (1996) Nerve growth factor: from neurotrophin to neurokine. *Trends Neurosci*, **19**, 514-520.

Li, H., Li, S. H., Yu, Z. X., Shelbourne, P. and Li, X. J. (2001) Huntington aggregate-associated axonal degeneration is an early pathological event in

- Huntington's disease mice. *J Neurosci*, **21**, 8473-8481.
- Li, H., Wyman, T., Yu, Z. X., Li, S. H. and Li, X. J. (2003a) Abnormal association of mutant huntingtin with synaptic vesicles inhibits glutamate release. *Hum Mol Genet*, **12**, 2021-2030.
- Li, J. Y., Leitner, B., Lovisetti-Scamihorn, P., Winkler, H. and Dahlstrom, A. (1999a) Proteolytic processing, axonal transport and differential distribution of chromogranins A and B, and secretogranin II (secretoneurin) in rat sciatic nerve and spinal cord. *Eur J Neurosci*, **11**, 528-544.
- Li, S., Finley, J., Liu, Z. J. et al. (2002a) Crystal structure of the cytoskeleton-associated protein glycine-rich (CAP-Gly) domain. *J Biol Chem*, **277**, 48596-48601.
- Li, S. H., Gutekunst, C. A., Hersch, S. M. and Li, X. J. (1998a) Association of HAP1 isoforms with a unique cytoplasmic structure. *J Neurochem*, **71**, 2178-2185.
- Li, S. H., Gutekunst, C. A., Hersch, S. M. and Li, X. J. (1998b) Interaction of huntingtin-associated protein with dynactin P150Glued. *J Neurosci*, **18**, 1261-1269.
- Li, S. H., Hosseini, S. H., Gutekunst, C. A., Hersch, S. M., Ferrante, R. J. and Li, X. J. (1998c) A human HAP1 homologue. Cloning, expression, and interaction with huntingtin. *J Biol Chem*, **273**, 19220-19227.
- Li, S. H., Yu, Z. X., Li, C. L., Nguyen, H. P., Zhou, Y. X., Deng, C. and Li, X.

- J. (2003b) Lack of huntingtin-associated protein-1 causes neuronal death resembling hypothalamic degeneration in Huntington's disease. *J Neurosci*, **23**, 6956-6964.
- Li, W. P., Xian, C., Rush, R. A. and Zhou, X. F. (1999b) Upregulation of brain-derived neurotrophic factor and neuropeptide Y in the dorsal ascending sensory pathway following sciatic nerve injury in rat. *Neurosci Lett*, **260**, 49-52.
- Li, X. J. and Li, S. H. (2005) HAP1 and intracellular trafficking. *Trends Pharmacol Sci*, **26**, 1-3.
- Li, X. J., Li, S. H., Sharp, A. H., Nucifora, F. C., Jr., Schilling, G., Lanahan, A., Worley, P., Snyder, S. H. and Ross, C. A. (1995) A huntingtin-associated protein enriched in brain with implications for pathology. *Nature*, **378**, 398-402.
- Li, X. J., Sharp, A. H., Li, S. H., Dawson, T. M., Snyder, S. H. and Ross, C. A. (1996) Huntingtin-associated protein (HAP1): discrete neuronal localizations in the brain resemble those of neuronal nitric oxide synthase. *Proc Natl Acad Sci U S A*, **93**, 4839-4844.
- Li, Y., Chin, L. S., Levey, A. I. and Li, L. (2002b) Huntingtin-associated protein 1 interacts with hepatocyte growth factor-regulated tyrosine kinase substrate and functions in endosomal trafficking. *J Biol Chem*, **277**, 28212-28221.

Reference

- Liu, Q. R., Lu, L., Zhu, X. G., Gong, J. P., Shaham, Y. and Uhl, G. R. (2006) Rodent BDNF genes, novel promoters, novel splice variants, and regulation by cocaine. *Brain Res*, **1067**, 1-12.
- Lou, H., Kim, S. K., Zaitsev, E., Snell, C. R., Lu, B. and Loh, Y. P. (2005) Sorting and activity-dependent secretion of BDNF require interaction of a specific motif with the sorting receptor carboxypeptidase e. *Neuron*, **45**, 245-255.
- Lu, B. (2003) Pro-region of neurotrophins: role in synaptic modulation. *Neuron*, **39**, 735-738.
- Lu, B. (2004) Acute and long-term synaptic modulation by neurotrophins. *Prog Brain Res*, **146**, 137-150.
- Luo, W., Wickramasinghe, S. R., Savitt, J. M., Griffin, J. W., Dawson, T. M. and Ginty, D. D. (2007) A hierarchical NGF signaling cascade controls Ret-dependent and Ret-independent events during development of nonpeptidergic DRG neurons. *Neuron*, **54**, 739-754.
- Maggirwar, S. B., Ramirez, S., Tong, N., Gelbard, H. A. and Dewhurst, S. (2000) Functional interplay between nuclear factor-kappaB and c-Jun integrated by coactivator p300 determines the survival of nerve growth factor-dependent PC12 cells. *J Neurochem*, **74**, 527-539.
- Maisonpierre, P. C., Belluscio, L., Friedman, B., Alderson, R. F., Wiegand, S. J., Furth, M. E., Lindsay, R. M. and Yancopoulos, G. D. (1990a) NT-3, BDNF,

- and NGF in the developing rat nervous system: parallel as well as reciprocal patterns of expression. *Neuron*, **5**, 501-509.
- Maisonpierre, P. C., Belluscio, L., Squinto, S., Ip, N. Y., Furth, M. E., Lindsay, R. M. and Yancopoulos, G. D. (1990b) Neurotrophin-3: a neurotrophic factor related to NGF and BDNF. *Science*, **247**, 1446-1451.
- Marco, S., Canudas, A. M., Canals, J. M., Gavalda, N., Perez-Navarro, E. and Alberch, J. (2002) Excitatory amino acids differentially regulate the expression of GDNF, neurturin, and their receptors in the adult rat striatum. *Exp Neurol*, **174**, 243-252.
- Martin, E. J., Kim, M., Velier, J. et al. (1999) Analysis of Huntington-associated protein 1 in mouse brain and immortalized striatal neurons. *J Comp Neurol*, **403**, 421-430.
- Matsudaira, P. (1991) Modular organization of actin crosslinking proteins. *Trends Biochem Sci*, **16**, 87-92.
- Mazella, J., Zsurger, N., Navarro, V. et al. (1998) The 100-kDa neurotensin receptor is gp95/sortilin, a non-G-protein-coupled receptor. *J Biol Chem*, **273**, 26273-26276.
- McBride, J. L., Ramaswamy, S., Gasmi, M. et al. (2006) Viral delivery of glial cell line-derived neurotrophic factor improves behavior and protects striatal neurons in a mouse model of Huntington's disease. *Proc Natl Acad Sci U S A*, **103**, 9345-9350.

Reference

- McDonald, N. Q., Lapatto, R., Murray-Rust, J., Gunning, J., Wlodawer, A. and Blundell, T. L. (1991) New protein fold revealed by a 2.3-A resolution crystal structure of nerve growth factor. *Nature*, **354**, 411-414.
- McGrail, M., Gepner, J., Silvanovich, A., Ludmann, S., Serr, M. and Hays, T. S. (1995) Regulation of cytoplasmic dynein function in vivo by the Drosophila Glued complex. *J Cell Biol*, **131**, 411-425.
- McGuire, J. R., Rong, J., Li, S. H. and Li, X. J. (2006) Interaction of Huntingtin-associated protein-1 with kinesin light chain: implications in intracellular trafficking in neurons. *J Biol Chem*, **281**, 3552-3559.
- Michalski, B. and Fahnestock, M. (2003) Pro-brain-derived neurotrophic factor is decreased in parietal cortex in Alzheimer's disease. *Brain Res Mol Brain Res*, **111**, 148-154.
- Middleton, G., Hamanoue, M., Enokido, Y., Wyatt, S., Pennica, D., Jaffray, E., Hay, R. T. and Davies, A. M. (2000) Cytokine-induced nuclear factor kappa B activation promotes the survival of developing neurons. *J Cell Biol*, **148**, 325-332.
- Milgram, S. L., Mains, R. E. and Eipper, B. A. (1996) Identification of routing determinants in the cytosolic domain of a secretory granule-associated integral membrane protein. *J Biol Chem*, **271**, 17526-17535.
- Minichiello, L., Calella, A. M., Medina, D. L., Bonhoeffer, T., Klein, R. and Korte, M. (2002) Mechanism of TrkB-mediated hippocampal long-term

- potentiation. *Neuron*, **36**, 121-137.
- Momose, Y., Murata, M., Kobayashi, K., Tachikawa, M., Nakabayashi, Y., Kanazawa, I. and Toda, T. (2002) Association studies of multiple candidate genes for Parkinson's disease using single nucleotide polymorphisms. *Ann Neurol*, **51**, 133-136.
- Moss, S. J. and Smart, T. G. (2001) Constructing inhibitory synapses. *Nat Rev Neurosci*, **2**, 240-250.
- Mowla, S. J., Farhadi, H. F., Pareek, S., Atwal, J. K., Morris, S. J., Seidah, N. G. and Murphy, R. A. (2001) Biosynthesis and post-translational processing of the precursor to brain-derived neurotrophic factor. *J Biol Chem*, **276**, 12660-12666.
- Mowla, S. J., Pareek, S., Farhadi, H. F., Petrecca, K., Fawcett, J. P., Seidah, N. G., Morris, S. J., Sossin, W. S. and Murphy, R. A. (1999) Differential sorting of nerve growth factor and brain-derived neurotrophic factor in hippocampal neurons. *J Neurosci*, **19**, 2069-2080.
- Mu, X., Silos-Santiago, I., Carroll, S. L. and Snider, W. D. (1993) Neurotrophin receptor genes are expressed in distinct patterns in developing dorsal root ganglia. *J Neurosci*, **13**, 4029-4041.
- Mukherjee, S., Ghosh, R. N. and Maxfield, F. R. (1997) Endocytosis. *Physiol Rev*, **77**, 759-803.
- Muragaki, Y., Chou, T. T., Kaplan, D. R., Trojanowski, J. Q. and Lee, V. M.

- (1997) Nerve growth factor induces apoptosis in human medulloblastoma cell lines that express TrkA receptors. *J Neurosci*, **17**, 530-542.
- Murer, M. G., Yan, Q. and Raisman-Vozari, R. (2001) Brain-derived neurotrophic factor in the control human brain, and in Alzheimer's disease and Parkinson's disease. *Prog Neurobiol*, **63**, 71-124.
- Murphy, R. A., Saide, J. D., Blanchard, M. H. and Young, M. (1977) Molecular properties of the nerve growth factor secreted in mouse saliva. *Proc Natl Acad Sci U S A*, **74**, 2672-2676.
- Murray, S. S. and Cheema, S. S. (2003) Constitutive expression of the low-affinity neurotrophin receptor and changes during axotomy-induced death of sensory neurones in the neonatal rat dorsal root ganglion. *J Anat*, **202**, 227-238.
- Nakata, T. and Hirokawa, N. (2003) Microtubules provide directional cues for polarized axonal transport through interaction with kinesin motor head. *J Cell Biol*, **162**, 1045-1055.
- Nangaku, M., Sato-Yoshitake, R., Okada, Y., Noda, Y., Takemura, R., Yamazaki, H. and Hirokawa, N. (1994) KIF1B, a novel microtubule plus end-directed monomeric motor protein for transport of mitochondria. *Cell*, **79**, 1209-1220.
- Nasir, J., Duan, K., Nichol, K. et al. (1998) Gene structure and map location of the murine homolog of the Huntington-associated protein, Hap1. *Mamm Genome*, **9**, 565-570.

Reference

- Neuwald, A. F., Aravind, L., Spouge, J. L. and Koonin, E. V. (1999) AAA+: A class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes. *Genome Res.*, **9**, 27-43.
- Nucifora, F. C., Jr., Sasaki, M., Peters, M. F. et al. (2001) Interference by huntingtin and atrophin-1 with cbp-mediated transcription leading to cellular toxicity. *Science*, **291**, 2423-2428.
- Numata, S., Ueno, S., Iga, J. et al. (2006) Brain-derived neurotrophic factor (BDNF) Val66Met polymorphism in schizophrenia is associated with age at onset and symptoms. *Neurosci Lett*, **401**, 1-5.
- Nykjaer, A., Lee, R., Teng, K. K. et al. (2004) Sortilin is essential for proNGF-induced neuronal cell death. *Nature*, **427**, 843-848.
- Page, K. J., Potter, L., Aronni, S., Everitt, B. J. and Dunnett, S. B. (1998) The expression of Huntington-associated protein (HAP1) mRNA in developing, adult and ageing rat CNS: implications for Huntington's disease neuropathology. *Eur J Neurosci*, **10**, 1835-1845.
- Pang, P. T., Teng, H. K., Zaitsev, E. et al. (2004) Cleavage of proBDNF by tPA/plasmin is essential for long-term hippocampal plasticity. *Science*, **306**, 487-491.
- Patapoutian, A. and Reichardt, L. F. (2001) Trk receptors: mediators of neurotrophin action. *Curr Opin Neurobiol*, **11**, 272-280.
- Qian, X. and Ginty, D. D. (2001) SH2-B and APS are multimeric adapters that

- augment TrkA signaling. *Mol Cell Biol*, **21**, 1613-1620.
- Quintyne, N. J., Gill, S. R., Eckley, D. M., Crego, C. L., Compton, D. A. and Schroer, T. A. (1999) Dynactin is required for microtubule anchoring at centrosomes. *J Cell Biol*, **147**, 321-334.
- Quintyne, N. J. and Schroer, T. A. (2002) Distinct cell cycle-dependent roles for dynein and dynein at centrosomes. *J Cell Biol*, **159**, 245-254.
- Radka, S. F., Holst, P. A., Fritzsche, M. and Altar, C. A. (1996) Presence of brain-derived neurotrophic factor in brain and human and rat but not mouse serum detected by a sensitive and specific immunoassay. *Brain Res*, **709**, 122-301.
- Rahman, A., Friedman, D. S. and Goldstein, L. S. (1998) Two kinesin light chain genes in mice. Identification and characterization of the encoded proteins. *J Biol Chem*, **273**, 15395-15403.
- Raiborg, C., Bache, K. G., Mehlum, A. and Stenmark, H. (2001) Function of Hrs in endocytic trafficking and signalling. *Biochem Soc Trans*, **29**, 472-475.
- Reichardt, L. F. (2006) Neurotrophin-regulated signalling pathways. *Philos Trans R Soc Lond B Biol Sci*, **361**, 1545-1564.
- Richardson, P. M., McGuinness, U. M. and Aguayo, A. J. (1980) Axons from CNS neurons regenerate into PNS grafts. *Nature*, **284**, 264-265.
- Ringstedt, T., Ibanez, C. F. and Nosrat, C. A. (1999) Role of brain-derived neurotrophic factor in target invasion in the gustatory system. *J Neurosci*, **19**,

3507-3518.

Rong, J., McGuire, J. R., Fang, Z. H., Sheng, G., Shin, J. Y., Li, S. H. and Li, X. J. (2006) Regulation of intracellular trafficking of huntingtin-associated protein-1 is critical for TrkA protein levels and neurite outgrowth. *J Neurosci*, **26**, 6019-6030.

Rybakowski, J. K., Borkowska, A., Czerski, P. M., Skibinska, M. and Hauser, J. (2003) Polymorphism of the brain-derived neurotrophic factor gene and performance on a cognitive prefrontal test in bipolar patients. *Bipolar Disord*, **5**, 468-472.

Sadri-Vakili, G. and Cha, J. H. (2006) Mechanisms of disease: Histone modifications in Huntington's disease. *Nat Clin Pract Neurol*, **2**, 330-338.

Sadri-Vakili, G., Menon, A. S., Farrell, L. A., Keller-McGandy, C. E., Cantuti-Castelvetro, I., Standaert, D. G., Augood, S. J., Yohrling, G. J. and Cha, J. H. (2006) Huntingtin inclusions do not down-regulate specific genes in the R6/2 Huntington's disease mouse. *Eur J Neurosci*, **23**, 3171-3175.

Samso, M., Radermacher, M., Frank, J. and Koonce, M. P. (1998) Structural characterization of a dynein motor domain. *J Mol Biol*, **276**, 927-937.

Saudou, F., Finkbeiner, S., Devys, D. and Greenberg, M. E. (1998) Huntingtin acts in the nucleus to induce apoptosis but death does not correlate with the formation of intranuclear inclusions. *Cell*, **95**, 55-66.

Schecterson, L. C. and Bothwell, M. (1992) Novel roles for neurotrophins are

- suggested by BDNF and NT-3 mRNA expression in developing neurons. *Neuron*, **9**, 449-463.
- Schinder, A. F., Berninger, B. and Poo, M. (2000) Postsynaptic target specificity of neurotrophin-induced presynaptic potentiation. *Neuron*, **25**, 151-163.
- Schliwa, M. and Woehlke, G. (2003) Molecular motors. *Nature*, **422**, 759-765.
- Schmidt, A. and Hall, A. (2002) Guanine nucleotide exchange factors for Rho GTPases: turning on the switch. *Genes Dev*, **16**, 1587-1609.
- Schnapp, B. J. (2003) Trafficking of signaling modules by kinesin motors. *J Cell Sci*, **116**, 2125-2135.
- Schnell, L., Schneider, R., Kolbeck, R., Barde, Y. A. and Schwab, M. E. (1994) Neurotrophin-3 enhances sprouting of corticospinal tract during development and after adult spinal cord lesion. *Nature*, **367**, 170-173.
- Schroer, T. A. (2004) Dynactin. *Annu Rev Cell Dev Biol*, **20**, 759-779.
- Seaman, M. N., McCaffery, J. M. and Emr, S. D. (1998) A membrane coat complex essential for endosome-to-Golgi retrograde transport in yeast. *J Cell Biol*, **142**, 665-681.
- Sedel, F., Bechade, C. and Triller, A. (1999) Nerve growth factor (NGF) induces motoneuron apoptosis in rat embryonic spinal cord in vitro. *Eur J Neurosci*, **11**, 3904-3912.
- Seidah, N. G., Benjannet, S., Pareek, S., Chretien, M. and Murphy, R. A.

- (1996) Cellular processing of the neurotrophin precursors of NT3 and BDNF by the mammalian proprotein convertases. *FEBS Lett*, **379**, 247-250.
- Selby, M. J., Edwards, R., Sharp, F. and Rutter, W. J. (1987) Mouse nerve growth factor gene: structure and expression. *Mol Cell Biol*, **7**, 3057-3064.
- Setou, M., Seog, D. H., Tanaka, Y., Kanai, Y., Takei, Y., Kawagishi, M. and Hirokawa, N. (2002) Glutamate-receptor-interacting protein GRIP1 directly steers kinesin to dendrites. *Nature*, **417**, 83-87.
- Shakiryanova, D., Tully, A., Hewes, R. S., Deitcher, D. L. and Levitan, E. S. (2005) Activity-dependent liberation of synaptic neuropeptide vesicles. *Nat Neurosci*, **8**, 173-178.
- Sharp, A. H., Loev, S. J., Schilling, G. et al. (1995) Widespread expression of Huntington's disease gene (IT15) protein product. *Neuron*, **14**, 1065-1074.
- Sheng, G., Chang, G. Q., Lin, J. Y. et al. (2006) Hypothalamic huntingtin-associated protein 1 as a mediator of feeding behavior. *Nat Med*, **12**, 526-533.
- Shimizu, E., Hashimoto, K. and Iyo, M. (2004) Ethnic difference of the BDNF 196G/A (val66met) polymorphism frequencies: the possibility to explain ethnic mental traits. *Am J Med Genet B Neuropsychiatr Genet*, **126**, 122-123.
- Sieghart, W. and Sperk, G. (2002) Subunit composition, distribution and function of GABA(A) receptor subtypes. *Curr Top Med Chem*, **2**, 795-816.
- Smeyne, R. J., Klein, R., Schnapp, A., Long, L. K., Bryant, S., Lewin, A., Lira, S. A. and Barbacid, M. (1994) Severe sensory and sympathetic neuropathies in

- mice carrying a disrupted Trk/NGF receptor gene. *Nature*, **368**, 246-249.
- Smith, M. A., Zhang, L. X., Lyons, W. E. and Mamounas, L. A. (1997) Anterograde transport of endogenous brain-derived neurotrophic factor in hippocampal mossy fibers. *Neuroreport*, **8**, 1829-1834.
- Smith, M. M., Shi, L. and Navre, M. (1995) Rapid identification of highly active and selective substrates for stromelysin and matrilysin using bacteriophage peptide display libraries. *J Biol Chem*, **270**, 6440-6449.
- Spires, T. L., Grote, H. E., Varshney, N. K., Cordery, P. M., van Dellen, A., Blakemore, C. and Hannan, A. J. (2004) Environmental enrichment rescues protein deficits in a mouse model of Huntington's disease, indicating a possible disease mechanism. *J Neurosci*, **24**, 2270-2276.
- Steffan, J. S., Kazantsev, A., Spasic-Boskovic, O. et al. (2000) The Huntington's disease protein interacts with p53 and CREB-binding protein and represses transcription. *Proc Natl Acad Sci U S A*, **97**, 6763-6768.
- Sugars, K. L. and Rubinsztein, D. C. (2003) Transcriptional abnormalities in Huntington disease. *Trends Genet*, **19**, 233-238.
- Sunderland, S. (1990) The anatomy and physiology of nerve injury. *Muscle Nerve*, **13**, 771-784.
- Sunderland, S. and Bradley, K. C. (1950) Endoneurial tube shrinkage in the distal segment of a severed nerve. *J Comp Neurol*, **93**, 411-420.
- Takeda, S., Yamazaki, H., Seog, D. H., Kanai, Y., Terada, S. and Hirokawa, N.

- (2000) Kinesin superfamily protein 3 (KIF3) motor transports fodrin-associating vesicles important for neurite building. *J Cell Biol*, **148**, 1255-1265.
- Tanaka, Y., Kanai, Y., Okada, Y., Nonaka, S., Takeda, S., Harada, A. and Hirokawa, N. (1998) Targeted disruption of mouse conventional kinesin heavy chain, kif5B, results in abnormal perinuclear clustering of mitochondria. *Cell*, **93**, 1147-1158.
- Tang, T. S., Tu, H., Chan, E. Y., Maximov, A., Wang, Z., Wellington, C. L., Hayden, M. R. and Bezprozvanny, I. (2003) Huntingtin and huntingtin-associated protein 1 influence neuronal calcium signaling mediated by inositol-(1,4,5) triphosphate receptor type 1. *Neuron*, **39**, 227-239.
- Tang, T. S., Tu, H., Orban, P. C., Chan, E. Y., Hayden, M. R. and Bezprozvanny, I. (2004) HAP1 facilitates effects of mutant huntingtin on inositol 1,4,5-trisphosphate-induced Ca release in primary culture of striatal medium spiny neurons. *Eur J Neurosci*, **20**, 1779-1787.
- Teng, H. K., Teng, K. K., Lee, R. et al. (2005) ProBDNF induces neuronal apoptosis via activation of a receptor complex of p75NTR and sortilin. *J Neurosci*, **25**, 5455-5463.
- Tessarollo, L., Tsoulfas, P., Martin-Zanca, D., Gilbert, D. J., Jenkins, N. A., Copeland, N. G. and Parada, L. F. (1993) trkB, a receptor for neurotrophin-3, is widely expressed in the developing nervous system and in non-neuronal tissues. *Development*, **118**, 463-475.

- Thomas, M. S., Zhang, W., Jordan, P. M., Saragovi, H. U. and Taglialatela, G. (2005) Signaling pathways mediating a selective induction of nitric oxide synthase II by tumor necrosis factor alpha in nerve growth factor-responsive cells. *J Neuroinflammation*, **2**, 19.
- Tonra, J. R., Curtis, R., Wong, V. et al. (1998) Axotomy upregulates the anterograde transport and expression of brain-derived neurotrophic factor by sensory neurons. *J Neurosci*, **18**, 4374-4383.
- Trushina, E., Dyer, R. B., Badger, J. D., 2nd et al. (2004) Mutant huntingtin impairs axonal trafficking in mammalian neurons in vivo and in vitro. *Mol Cell Biol*, **24**, 8195-8209.
- Utley, D. S., Lewin, S. L., Cheng, E. T., Verity, A. N., Sierra, D. and Terris, D. J. (1996) Brain-derived neurotrophic factor and collagen tubulization enhance functional recovery after peripheral nerve transection and repair. *Arch Otolaryngol Head Neck Surg*, **122**, 407-413.
- Vallee, R. B., Williams, J. C., Varma, D. and Barnhart, L. E. (2004) Dynein: An ancient motor protein involved in multiple modes of transport. *J Neurobiol*, **58**, 189-200.
- Vanhaesebroeck, B., Leevers, S. J., Ahmadi, K., Timms, J., Katso, R., Driscoll, P. C., Woscholski, R., Parker, P. J. and Waterfield, M. D. (2001) Synthesis and function of 3-phosphorylated inositol lipids. *Annu Rev Biochem*, **70**, 535-602.
- Vaughan, P. S., Miura, P., Henderson, M., Byrne, B. and Vaughan, K. T.

- (2002) A role for regulated binding of p150(Glued) to microtubule plus ends in organelle transport. *J Cell Biol*, **158**, 305-319.
- Volosin, M., Song, W., Almeida, R. D., Kaplan, D. R., Hempstead, B. L. and Friedman, W. J. (2006) Interaction of survival and death signaling in basal forebrain neurons: roles of neurotrophins and proneurotrophins. *J Neurosci*, **26**, 7756-7766.
- von Bartheld, C. S. (2004) Axonal transport and neuronal transcytosis of trophic factors, tracers, and pathogens. *J Neurobiol*, **58**, 295-314.
- von Bartheld, C. S., Byers, M. R., Williams, R. and Bothwell, M. (1996a) Anterograde transport of neurotrophins and axodendritic transfer in the developing visual system. *Nature*, **379**, 830-833.
- von Bartheld, C. S., Williams, R., Lefcort, F., Clary, D. O., Reichardt, L. F. and Bothwell, M. (1996b) Retrograde transport of neurotrophins from the eye to the brain in chick embryos: roles of the p75NTR and trkB receptors. *J Neurosci*, **16**, 2995-3008.
- Wahle, T., Prager, K., Raffler, N., Haass, C., Famulok, M. and Walter, J. (2005) GGA proteins regulate retrograde transport of BACE1 from endosomes to the trans-Golgi network. *Mol Cell Neurosci*, **29**, 453-461.
- Wang, H., Wu, L. L., Song, X. Y., Luo, X. G., Zhong, J. H., Rush, R. A. and Zhou, X. F. (2006) Axonal transport of BDNF precursor in primary sensory neurons. *Eur J Neurosci*, **24**, 2444-2452.

- Wang, K. C., Kim, J. A., Sivasankaran, R., Segal, R. and He, Z. (2002) P75 interacts with the Nogo receptor as a co-receptor for Nogo, MAG and OMgp. *Nature*, **420**, 74-78.
- Waterman-Storer, C. M., Karki, S. and Holzbaur, E. L. (1995) The p150Glued component of the dynein complex binds to both microtubules and the actin-related protein centrin (Arp-1). *Proc Natl Acad Sci U S A*, **92**, 1634-1638.
- Waterman-Storer, C. M., Karki, S. B., Kuznetsov, S. A., Tabb, J. S., Weiss, D. G., Langford, G. M. and Holzbaur, E. L. (1997) The interaction between cytoplasmic dynein and dynein is required for fast axonal transport. *Proc Natl Acad Sci U S A*, **94**, 12180-12185.
- Weibezahn, J., Bukau, B. and Mogk, A. (2004) Unscrambling an egg: protein disaggregation by AAA+ proteins. *Microb Cell Fact*, **3**, 1.
- Wiesmann, C., Ultsch, M. H., Bass, S. H. and de Vos, A. M. (1999) Crystal structure of nerve growth factor in complex with the ligand-binding domain of the TrkA receptor. *Nature*, **401**, 184-188.
- Williams, R., Gaber, B. and Gunning, J. (1982) Raman spectroscopic determination of the secondary structure of crystalline nerve growth factor. *J Biol Chem*, **257**, 13321-13323.
- Wong, R. W., Setou, M., Teng, J., Takei, Y. and Hirokawa, N. (2002a) Overexpression of motor protein KIF17 enhances spatial and working memory in transgenic mice. *Proc Natl Acad Sci U S A*, **99**, 14500-14505.

Reference

- Wong, S. T., Henley, J. R., Kanning, K. C., Huang, K. H., Bothwell, M. and Poo, M. M. (2002b) A p75(NTR) and Nogo receptor complex mediates repulsive signaling by myelin-associated glycoprotein. *Nat Neurosci*, **5**, 1302-1308.
- Woo, N. H., Teng, H. K., Siao, C. J., Chiaruttini, C., Pang, P. T., Milner, T. A., Hempstead, B. L. and Lu, B. (2005) Activation of p75NTR by proBDNF facilitates hippocampal long-term depression. *Nat Neurosci*, **8**, 1069-1077.
- Wooten, M. W., Seibenhener, M. L., Zhou, G., Vandenplas, M. L. and Tan, T. H. (1999) Overexpression of atypical PKC in PC12 cells enhances NGF-responsiveness and survival through an NF-kappaB dependent pathway. *Cell Death Differ*, **6**, 753-764.
- Xia, C., Rahman, A., Yang, Z. and Goldstein, L. S. (1998) Chromosomal localization reveals three kinesin heavy chain genes in mouse. *Genomics*, **52**, 209-213.
- Xing, J., Kornhauser, J. M., Xia, Z., Thiele, E. A. and Greenberg, M. E. (1998) Nerve growth factor activates extracellular signal-regulated kinase and p38 mitogen-activated protein kinase pathways to stimulate CREB serine 133 phosphorylation. *Mol Cell Biol*, **18**, 1946-1955.
- Xu, X. M., Guenard, V., Kleitman, N., Aebsicher, P. and Bunge, M. B. (1995) A combination of BDNF and NT-3 promotes supraspinal axonal regeneration into Schwann cell grafts in adult rat thoracic spinal cord. *Exp Neurol*, **134**, 261-

272.

Yamada, K. and Nabeshima, T. (2003) Brain-derived neurotrophic factor/TrkB

signaling in memory processes. *J Pharmacol Sci*, **91**, 267-270.

Yamashita, T., Fujitani, M., Hata, K., Mimura, F. and Yamagishi, S. (2005)

Diverse functions of the p75 neurotrophin receptor. *Anat Sci Int*, **80**, 37-41.

Yamashita, T. and Tohyama, M. (2003) The p75 receptor acts as a displacement factor that releases Rho from Rho-GDI. *Nat Neurosci*, **6**, 461-

467.

Yamashita, T., Tucker, K. L. and Barde, Y. A. (1999) Neurotrophin binding to

the p75 receptor modulates Rho activity and axonal outgrowth. *Neuron*, **24**,

585-593.

Yano, H. and Chao, M. V. (2004) Mechanisms of neurotrophin receptor

vesicular transport. *J Neurobiol*, **58**, 244-257.

Yano, H., Lee, F. S., Kong, H., Chuang, J., Arevalo, J., Perez, P., Sung, C. and

Chao, M. V. (2001) Association of Trk neurotrophin receptors with

components of the cytoplasmic dynein motor. *J Neurosci*, **21**, RC125.

Ye, J. H. and Houle, J. D. (1997) Treatment of the chronically injured spinal

cord with neurotrophic factors can promote axonal regeneration from

supraspinal neurons. *Exp Neurol*, **143**, 70-81.

Yeiser, E. C., Rutkoski, N. J., Naito, A., Inoue, J. and Carter, B. D. (2004)

Neurotrophin signaling through the p75 receptor is deficient in *traf6*-/- mice. *J*

- Neurosci*, **24**, 10521-10529.
- Yin, Q., Kemp, G. J. and Frostick, S. P. (1998) Neurotrophins, neurones and peripheral nerve regeneration. *J Hand Surg [Br]*, **23**, 433-437.
- Yonekawa, Y., Harada, A., Okada, Y., Funakoshi, T., Kanai, Y., Takei, Y., Terada, S., Noda, T. and Hirokawa, N. (1998) Defect in synaptic vesicle precursor transport and neuronal cell death in KIF1A motor protein-deficient mice. *J Cell Biol*, **141**, 431-441.
- Yoon, S. O., Casaccia-Bonelli, P., Carter, B. and Chao, M. V. (1998) Competitive signaling between TrkB and p75 nerve growth factor receptors determines cell survival. *J Neurosci*, **18**, 3273-3281.
- Yuan, J. and Yankner, B. A. (2000) Apoptosis in the nervous system. *Nature*, **407**, 802-809.
- Yuan, X. B., Jin, M., Xu, X., Song, Y. Q., Wu, C. P., Poo, M. M. and Duan, S. (2003) Signalling and crosstalk of Rho GTPases in mediating axon guidance. *Nat Cell Biol*, **5**, 38-45.
- Zampieri, N. and Chao, M. V. (2006) Mechanisms of neurotrophin receptor signalling. *Biochem Soc Trans*, **34**, 607-611.
- Zeron, M. M., Hansson, O., Chen, N., Wellington, C. L., Leavitt, B. R., Brundin, P., Hayden, M. R. and Raymond, L. A. (2002) Increased sensitivity to N-methyl-D-aspartate receptor-mediated excitotoxicity in a mouse model of Huntington's disease. *Neuron*, **33**, 849-860.

Reference

- Zhao, C., Takita, J., Tanaka, Y. et al. (2001) Charcot-Marie-Tooth disease type 2A caused by mutation in a microtubule motor KIF1Bbeta. *Cell*, **105**, 587-597.
- Zhou, X. F., Chie, E. T., Deng, Y. S., Zhong, J. H., Xue, Q., Rush, R. A. and Xian, C. J. (1999) Injured primary sensory neurons switch phenotype for brain-derived neurotrophic factor in the rat. *Neuroscience*, **92**, 841-853.
- Zhou, X. F., Li, W. P., Zhou, F. H., Zhong, J. H., Mi, J. X., Wu, L. L. and Xian, C. J. (2005) Differential effects of endogenous brain-derived neurotrophic factor on the survival of axotomized sensory neurons in dorsal root ganglia: a possible role for the p75 neurotrophin receptor. *Neuroscience*, **132**, 591-603.
- Zhou, X. F. and Rush, R. A. (1996) Endogenous brain-derived neurotrophic factor is anterogradely transported in primary sensory neurons. *Neuroscience*, **74**, 945-953.
- Zhou, X. F., Song, X. Y., Zhong, J. H., Barati, S., Zhou, F. H. and Johnson, S. M. (2004) Distribution and localization of pro-brain-derived neurotrophic factor-like immunoreactivity in the peripheral and central nervous system of the adult rat. *J Neurochem*, **91**, 704-715.
- Zuccato, C., Ciampola, A., Rigamonti, D. et al. (2001) Loss of huntingtin-mediated BDNF gene transcription in Huntington's disease. *Science*, **293**, 493-498.
- Zuccato, C., Tartari, M., Crotti, A. et al. (2003) Huntingtin interacts with

Reference

REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes.

Nat Genet, **35**, 76-83.

Zucker, B., Luthi-Carter, R., Kama, J. A., Dunah, A. W., Stern, E. A., Fox, J. H., Standaert, D. G., Young, A. B. and Augood, S. J. (2005) Transcriptional dysregulation in striatal projection- and interneurons in a mouse model of Huntington's disease: neuronal selectivity and potential neuroprotective role of HAP1. *Hum Mol Genet*, **14**, 179-189.