

**REFERENCES:**

- Anders, F.J. & Leatherman, S.P. (1987). Disturbance of beach sediment by off-road vehicles. Environmental Geology and Water Sciences **9**: 183 – 189.
- Anderson, M.J. (2006). Distance-based tests for homogeneity of multivariate dispersions. Biometrics **62**: 245 – 253.
- Anderson, M.J., Gorley, R.N. & Clarke, K.R. (2008). PERMANOVA+ for PRIMER: Guide to software and statistical methods. PRIMER-E: Plymouth, UK.
- Anonymous (2008). Vehicles on beaches final report to the City of Onkaparinga October 2008. Flinders Research Centre for Coastal and Catchment Environments, Flinders University, Adelaide, South Australia.
- Armonies, W. & Reise, K. (2000). Faunal diversity across a sandy shore. Marine Ecology-Progress Series **196**: 49 – 57.
- Arthington, A.H., Yeates, G.W. & Conrick, D.L. (1986). Nematodes, including a new record of *Tobrillus diversipapillatus* in Australia, as potential indicators of sewage effluent pollution. Australian Journal of Marine and Freshwater Research **37**: 159 – 166.
- Aubrey, D.G. (1983). Beach changes on coasts with different wave climates. In: McLachlan, A. & Th Erasmus (eds.) *Sandy beaches as ecosystems* Dr. W. Junk Publishers. The Hague. 63 – 85.
- Baker-Gabb, D. & Weston, M. (2006). South Australian recovery plan for the hooded plover, *Thinornis rubricollis*. Appendix 14 in Department for Environment and Heritage Hooded Plover recovery plan, Southern Fleurieu Coastal Action Plan and Conservation Priority Study.
- Bale, J.J. & Kenny, A.J. (2005). Sediment analysis and seabed characterisation. In: *Methods for the study of the marine benthos*. A. Eleftheriou and A. McIntyre (eds.). Blackwell Science Ltd. New Delhi, India. pp. 43 – 86.
- Barros, F. (2001). Ghost crabs as a tool for rapid assessment of human impacts on exposed sandy beaches. Biological Conservation **97**: 399 – 404.

- Bartsch, I. (2006). Halacaroidea (Acari): a guide to marine genera. Organisms Diversity and Evolution **6S**: 1 – 104.
- Bell, T.-A. (2005). The impact of four-wheel drive vehicles on, and the natural variation of, sandy beach meiofauna. Masters Thesis, University of New England.
- Blott, S.J. & Pye, K. (2001). GRADISTAT: A grain size distribution and statistics package for the analysis of unconsolidated sediments. Earth Surface Processes and Landforms **26**: 1237 – 1248.
- Blott, S.J. & Pye, K. (2006). Particle size distribution analysis of sand-sized particles by laser diffraction: an experimental investigation of instrument sensitivity and the effects of particle shape. Sedimentology **53**: 671 – 685.
- Bodin, P. & Lemoal, Y. (1982). Short-dated effects, in meiofaunal and macrofauna, of the cleaning of a beach polluted by hydrocarbons with utilization of an oil-spill dispersant. Acta Oecologica-Oecologia Applicata **3**: 263 – 280.
- Bongers, T. (1990). The maturity index: an ecological measure of environmental disturbance based on nematode species composition. Oecologia **83**: 14 – 19.
- Bongers, T. & Bongers, M. (1998). Functional diversity of nematodes. Applied Soil Ecology **10**: 239 – 251.
- Bongers, T., Alkemade, R. & Yeates, G.W. (1991). Interpretation of disturbance-induced maturity decrease in marine nematode assemblages by means of the maturity index. Marine Ecology-Progress Series **76**: 135 – 142.
- Brazeiro, A. & Defeo, O. (1996). Macroinfauna zonation in microtidal sandy beaches: is it possible to identify patterns in such variable environments? Estuarine, Coastal and Shelf Science **42**: 523 – 536.
- Brazeiro, A. & Defeo, O. (1999). Effects of harvesting and density-dependence on the demography of sandy beach populations: the yellow clam *Mesodesmata mactroides* of Uruguay. Marine Ecology-Progress Series **182**: 127 – 135.

- Brown, A.C. & McLachlan, A. (2002). Sandy shore ecosystems and the threats facing them: some predictions for the year 2025. Environmental Conservation **29**: 62 – 77.
- Caton, B. (2007). The impact of climate change on the coastal lands of the City of Onkaparinga. A report for the City of Onkaparinga 18 July 2007.
- Chapman, M.G., Underwood, A.J. & Skilleter, G.A. (1995). Variability at different spatial scales between a subtidal assemblage exposed to the discharge of sewage at two control locations. *Journal of Experimental Marine Biology and Ecology* **189**: 103 – 122.
- Clarke, K.R. & Green, R.H. (1988). Statistical design and analysis for a 'biological effects' study. *Marine Ecology-Progress Series* **46**: 213 – 226.
- Clarke, K.R. & Gorley, R.N. (2006). PRIMER v6: User manual/tutorial. PRIMER-E: Plymouth, UK.
- Clarke, K.R. & Warwick, R.M. (2001). Change in marine communities, 2<sup>nd</sup> Edition. PRIMER-E Ltd., Plymouth.
- Colombini, I. & Chelazzi, L. (2003). Influence of marine allochthonous input on sandy beach communities. Oceanography and Marine Biology **41**: 115 – 159.
- Colombini, I., Fallaci, M., Milanese, F., Scapini, F. & Chelazzi, L. (2003). Comparative diversity in sandy littoral ecosystems of the western Mediterranean. Estuarine, Coastal and Shelf Science **58S**: 93 – 104.
- Dahl, E. (1952). Some aspects of the ecology and zonation of the fauna on sandy beaches. Oikos **4**: 1 – 27.
- Davies, V.T. (1986). Australian spiders (Araneae). Collection, preservation and identification. Queensland Museum, Brisbane.
- Davis, R.A. Jr. & Fox, W.T. (1972). Coastal process and nearshore sand bars. Journal of Sedimentary Petrology **42**: 401 – 412.
- De Ley, P., Decraemer, W. & Abebe, E. (2006). Introduction: summary of present knowledge and research addressing the ecology and taxonomy of freshwater nematodes. In: *Freshwater nematodes, ecology and taxonomy*. E. Abebe, I. Andrassy & W. Transpurger (eds.), CABI Publishing, Wallingford, UK. pp. 3 – 30.

- Dean, R.G. (1973). Heuristic models of sand transport in the surf zone. Proc. Conf. Eng. Dyn. Surf Zone, American Society of Civil Engineers. New York.
- Defeo, O. & McLachlan, A. (2005). Patterns, processes and regulatory mechanism in sandy beach macrofauna: a multi-scale analysis. Marine Ecology-Progress Series **295**: 1 – 20.
- Defeo, O. & Rueda, M. (2002). Spatial structure, sampling design and abundance estimates in sandy beach macroinfauna: some warnings and new perspectives. Marine Biology **140**: 1215 – 1225.
- Defeo, O., Jaramillo, E. & Lyonnet, A. (1992). Community structure and intertidal zonation of the macroinfauna in the Atlantic coast of Uruguay. Journal of Coastal Research **8**: 830 – 839.
- Defeo, O., McLachlan, A., Schoeman, D.S., Schlacher, T.A., Dugan, J., Jones, A., Lastra, M. & Scapini, F. (2009). Threats to sandy beach ecosystems: a review. Estuarine, Coastal and Shelf Science **81**: 1 – 12.
- Di Domenico, M., da Cunha Lana, P. & Garraffoni, A.R.S. (2009). Distribution patterns of interstitial polychaetes in sandy beaches of southern Brazil. Marine Ecology **30**: 47 – 62.
- Dole-Olivier, M.-J., Galassi, D.M.P., Marmonier, P. & Creuzé des Châtelliers, M. (2000). The biology and ecology of lotic microcrustaceans. Freshwater Biology **44**: 63 – 91.
- Downes, B.J., Barmuta, L.A., Fairweather, P.G., Faith, D.P., Keough, Lake, P.S., Mapstone, B. & Quinn, G.P. (2002). *Monitoring ecological impacts: concepts and practice in flowing waters*. Cambridge University Press, Cambridge.
- Dugan, J.E., Hubbard, D.M., McCrary, M.D. & Peirson, M.O. (2003). The response of macrofauna communities and shorebirds to macrophyte wrack subsidies on exposed sandy beaches of southern California. Estuarine, Coastal and Shelf Science **58S**: 25 – 40.
- Dugan, J.E., Hubbard, D.M., Rodil, I.F., Revell, D.L. & Schroeter, S. (2008). Ecological effects of coastal armouring on sandy beaches. Marine Ecology **29S1**: 160 – 170.
- Dugan, J.E., Page, H.M., Hubbard, D., Schooler, N. & Schroeter, S. (2009). Ecological recovery of disturbed beach ecosystems. *In* 5<sup>th</sup> International

- Symposium on Sandy Beaches (Abstracts). University of Mohammed V Agdal, Scientific Institute, Rabat, Morocco, 19<sup>th</sup> – 23<sup>rd</sup> October, 2009.
- Duong, H.L.S. (2008). Investigating the ecological implications of wrack removal on South Australian sandy beaches. Unpublished PhD thesis. Flinders University, Adelaide, South Australia.
- Edgar, G.J. (2000). Australian marine life – the plants and animals of temperate waters. Revised Edition. Reed New Holland Pty. Ltd., Sydney.
- Ellis, M. (2006). Shorebird distributions and behavioural associations with beach wrack. Honours Thesis, Flinders University, Adelaide, South Australia.
- Fairbridge, R.W. (2004). Classification of coasts. Journal of Coastal Research **20**: 155 – 165.
- Fanini, L., Marchetti, G.M., Scapini, F. & Defeo, O. (2009). Effects of beach nourishment and groynes building on population and community descriptors of mobile arthropodofauna. Ecological Indicators **9**: 167 – 178.
- Fenchel, T.M. (1978). The ecology of the micro- and meiobenthos. Annual Review of Ecology and Systematics **9**: 99 – 121.
- Foster-Smith, J., Birchenough, A.C., Evans, S.M. & Prince, J. (2007). Human impacts on Cable Beach, Broome (Western Australia). Coastal Management **35**: 181 – 194.
- Gheskiere, T., Vincx, M., Greet, P. & Steven, D. (2006). Are strandline meiofauna assemblages affected by a once-only mechanical beach cleaning? Experimental findings. Marine Environmental Research **61**: 245 – 264.
- Gheskiere, T., Vincx, M., Weslawski, J.M., Scarpini, F. & Degraer, S. (2005). Meiofauna as descriptor of tourism-induced changes at sandy beaches. Marine Environmental Research **60**: 245 – 265.
- Godfrey, P.J. & Godfrey, M.M. (1981). Ecological effects of off-road vehicles on Cape Cod. Oceanus **23**: 56 – 67.
- Gooderham, J. & Tsyrlin, E. (2003). The waterbug book. CSIRO Publishing, Collingwood.
- Gowlett-Holmes, K. (2008). A field guide to the marine invertebrates of South Australia. Notomares, Sandy Bay.

- Gray, J.S. & Elliott, M. (2009). Ecology of marine sediments: from science to management. Second Edition. Oxford University Press, Oxford.
- Hagerman, G.M. Jr. & Reiger, R.M. (1981). Dispersal of benthic meiofauna by wave and current action in Brogue Sound, North Carolina, U.S.A. P.S.Z.N.I. Marine Ecology **2**: 245 – 270.
- Hale, H.M. (1927-29). The crustaceans of South Australia. Harrison Weir, Government Printer, Adelaide.
- Harriague, A.C., Petrillo, M., Della Croce, N., Panciroli, H., Pusceddu, A., Pelosi, G., Danovaro, R. & Albertelli, G. (2008). Intertidal benthic communities of two Chilean coastal islands (Santa Maria and Mocha, Southeastern Pacific). Chemistry and Ecology **24**: 119 – 128.
- Haynes, D. & Quinn, G.P. (1995). Temporal and spatial variability in community structure of a sandy intertidal beach, Cape Paterson, Victoria, Australia. Marine and Freshwater Research **46**: 931 – 942.
- Haynes, M.O. & Boothroyd, J.C. (1969). Storms as modifying agents in the coastal environment. In: M.O. Hayes (ed.), *Coastal Environments: NE Massachusetts*, Department of Geology, University of Massachusetts, Amhurst, pp. 290 – 315.
- Heck, K.L., Carruthers, T.J.B., Duarte, C.M., Hughes, A.R., Kendrick, G., Orth, R.J. & Williams, S.W. (2008). Trophic transfers from seagrass meadows subsidize diverse marine and terrestrial consumers. Ecosystems **11**: 1198 – 1210.
- Herrmann, M., Barreira, C.D.A.R., Arntz, W.E., Laudien, J. & Penchaszadeh, P.E. (2010). Testing the habitat harshness hypothesis: reproductive biology of the wedge clam *Donax hanleyanus* (Bivalvia: Donacidae) on three Argentinean sandy beaches with contrasting morphodynamics. Journal of Molluscan Studies **76**: 33 – 47.
- Hesp, P.A. (1999). The beach backshore and beyond. In: *Handbook of Beach and Shoreface Morphodynamics*, A.D. Short, (ed.), Wiley, London, pp. 145 – 170.
- Heymans, J.J. & McLachlan, A. (1996). Carbon budget and network analysis of a high-energy beach/surf-zone ecosystem. Estuarine, Coastal and Shelf Science **43**: 485 – 505.
- Higgins, R.P. & Thiel, H. (1988). Introduction to the study of meiofauna. Smithsonian Institution Press, Washington D.C.

- Hosier, P.E. & Eaton, T.E. (1980). The impact of vehicles on dune and grassland vegetation on a southeastern North Carolina barrier beach. Journal of Applied Ecology **17**: 173 – 183.
- Hosier, P.E., Kochhar, M. & Thayer, V. (1981). Off-road vehicle and pedestrian effects on the sea-approach of hatchling loggerhead turtles. Environmental Conservation **8**: 158 – 161.
- Hurlbert, S.H. (1984). Pseudoreplication and the design of ecological field experiments. Ecological Monographs **54**: 187 – 211.
- Ince, R., Hyndes, G.A., Lavery, P. & Vanderklift, M.A. (2007). Marine macrophytes directly enhance abundances of sandy beach fauna through provision of food and habitat. Estuarine, Coastal and Shelf Science **74**: 77 – 86.
- Inglis, G. (1989). The colonisation and degradation of stranded *Macrocystis pyrifera* (L.) C. Ag. By the macrofauna of a New Zealand sandy beach. Journal of Experimental Marine Biology and Ecology **125**: 203 – 217.
- James, R.J. (2000). From beaches to beach environments: linking the ecology, human-use and management of beaches in Australia. Ocean and Coastal Management **43**: 495 – 514.
- James, R.J. & Fairweather, P.G. (1996). Spatial variation of intertidal macrofauna on a sandy ocean beach in Australia. Estuarine, Coastal and Shelf Science **43**: 81 – 107.
- Jaramillo, E., McLachlan, A. & Coetzee, P. (1993). Intertidal zonation patterns of macroinfauna over a range of exposed sandy beaches in south-central Chile. Marine Ecology-Progress Series **101**: 105 – 118.
- Jensen, P. (1987). Feeding ecology of free-living aquatic nematodes. Marine Ecology-Progress Series **35**: 187 – 196.
- Kennedy, A. & Jacoby, C.A. (1999). Biological indicators of marine environmental health: meiofauna – a neglected benthic component? Environmental Monitoring and Assessment **54**: 47 – 68.
- Kirkman, H. & Kendrick, G.A. (1997). Ecological significance and commercial harvesting of drifting and beach-cast macro-algae and seagrasses in Australia: a review. Journal of Applied Phycology **9**: 311 – 326.
- Komar, P.D. (1976). Beach processes and sedimentation. Prentice-Hall Inc., New Jersey.

- Koop, K. & Griffiths, C.L. (1982). The relative significance of bacteria, meio- and macrofauna on an exposed sandy beach. Marine Biology **66**: 295 – 300.
- Kutiel, P., Eden, E. & Zhevelev, Y. (2000). Effect of experimental trampling and off-road motorcycle traffic on soil and vegetation of stabilized coastal dunes, Israel. Environmental Conservation **27**: 14 – 23.
- Leatherman, S.P. (1987). Field measurement in microtopography. Journal of Coastal Research **3**: 233 – 235.
- Leatherman, S.P. & Godfrey, P.J. (1979). The impact of off-road vehicles on coastal ecosystems in Cape Cod national seashore: An overview. UM/NPSCRU report no. 34. Amherst, Mass.: The Environmental Institute, Univ. Massachusetts.
- Leber, K.M. (1982). Seasonality of macroinvertebrates on a temperate, high wave energy sandy beach. Bulletin of Marine Science **32**: 86 – 98.
- Lecari, D. & Defeo, O. (2003). Variation of a sandy beach macroinvertebrate community along a human-induced environmental gradient. Estuarine, Coastal and Shelf Science **58S**: 17 – 24.
- Liddle, M.J. & Greig-Smith, P. (1975). A survey of tracks and paths in a sand dune ecosystem. I. Soils. Journal of Applied Ecology **12**: 893 – 908.
- Liddle, M.J. & Moore, K.G. (1974). The microclimate of sand dune tracks: the relative contribution of vegetation removal and soil compression. Journal of Applied Ecology **11**: 1057 – 1068.
- Lorenzen, S. Prein, M. & Valentin, C. (1987). Mass aggregations of the free living marine nematode *Pontonema vulgare* (Oncholaimidae) in organically polluted fjords, Marine Ecology-Progress Series **37**: 27 – 34.
- Luckenbach, R.A. & Bury, B. (1983). Effects of off-road vehicles on the biota of the Algodones dunes, Imperial County, California. Journal of Applied Ecology **20**: 265 – 286.
- Lucrezi, S., Schlacher, T.A. & Robinson, W. (2009). Human disturbance as a cause of bias in ecological indicators for sandy beaches: Experimental evidence for the effects of human trampling on ghost crabs (*Ocypode* spp.). Ecological Indicators **9**: 913 – 921.
- Malm, T., Raberg, S. Fell, S. & Carlsson, P. (2004). Effects of beach cast cleaning on beach quality, microbial food web, and littoral macrofaunal biodiversity. Estuarine, Coastal and Shelf Science **60**: 339 – 347.

- Mare, M.F. (1942). A study of a marine benthic community with special reference to the micro-organisms. Journal of the Marine Biological Association of the United Kingdom **25**: 517 – 863.
- Masselink, G. & Hughes, M.G. (2003). Introduction to coastal processes and geomorphology. Arnold, London.
- Masselink, G. & Pattiaratchi, C. B. (2001). Seasonal changes in beach morphology along the sheltered coastline of Perth, Western Australia. Marine Geology **172**: 243 – 263.
- Masselink, G. & Short, A.D. (1993). The effect of tide range on beach morphodynamics and morphology: a conceptual beach model. Journal of Coastal Research **9**: 785 – 800.
- Masselink, G., Auger, N., Russell, P. & O'Hare, T. (2007). Short-term morphological change and sediment dynamics in the intertidal zone of a macrotidal beach. Sedimentology **54**: 39 – 53.
- Masselink, G., Hegge, B.J. & Pattiaratchi, C.B. (1997). Beach cusp morphodynamics. Earth Processes and Landforms **22**: 1139 – 1155.
- Matthews, E.G. & Queale, L.F. (1997). Littoral Insects (Phylum Arthropoda: Class Insecta). Chapter 22 in: Marine invertebrates of southern Australia Part III. S.A. Shepherd & M. Davies (eds.). South Australian Research and Development Institute, Adelaide.
- Mazumder, D. & Saintilan, N. (2003). A comparison of sampling techniques in the assessment of burrowing crab abundance in saltmarsh and mangrove in temperate Australia. Wetlands (Aust) **21**: 1 – 15.
- McDonald, R. C., Isbell, R. F., Speight, J. G., Walker, J. & Hopkins, M. S. (1990). Australian soil and land survey field handbook, 2<sup>nd</sup> edition, Inkata Press Pty. Ltd., Melbourne and Sydney, Australia.
- McKechnie, J.M. (2003). Wrack: a nuisance or a vital component of sandy beach ecosystems? Unpublished Honours thesis, Flinders University, Adelaide, South Australia.
- McKechnie, J.M. & Fairweather, P.G. (2003). Ecological implications for the management of wrack on South Australian sandy beaches: A report to the Coast Protection Board. School of Biological Sciences, Flinders University, Adelaide, South Australia.

- McLachlan, A. (1979). Volumes of seawater filtered by East Cape sandy beaches. South African Journal of Science **75**: 75 – 79.
- McLachlan, A. (1980). The definition of sandy beaches in relation to exposure: a simple rating system. South African Journal of Science **76**: 137 – 138.
- McLachlan, A. (1985). The biomass of macro- and interstitial fauna on clean and wrack-covered beaches in Western Australia. Estuarine, Coastal and Shelf Science **21**: 587 – 599.
- McLachlan, A. (1989). Water filtration by dissipative beaches. Limnology and Oceanography **34**: 774 – 780.
- McLachlan, A. (1990). Dissipative beaches and macroinfaunal communities on exposed intertidal sands. Journal of Coastal Research **6**: 57 – 71.
- McLachlan, A. & Brown, A.C. (2006). Ecology of sandy shores. Elsevier Science, Amsterdam.
- McLachlan, A. & Dorvlo, A. (2005). Global patterns in sandy beach macrobenthic communities. Journal of Coastal Research **21**: 674 – 687.
- McLachlan, A. & Dorvlo, A. (2007). Global patterns in sandy beach macrobenthic communities: biological factors. Journal of Coastal Research **23**: 1081 – 1087.
- McLachlan, A. & Jaramillo, E. (1995). Zonation on sandy beaches. Oceanography and Marine Biology - an Annual Review **33**: 305 – 335.
- McLachlan, A., Cockcroft, A.C. & Malan, D.E. (1984). Benthic faunal responses to a high energy gradient. Marine Ecology-Progress Series **16**: 51 – 63.
- McLachlan, A., Eliot, I.G. & Clark, D.J. (1985). Water filtration through reflective microtidal beaches and shallow sublittoral sands and its implications for an inshore ecosystem in Western Australia. Estuarine, Coastal and Shelf Science **21**: 91 – 104.
- Moffett, M.D., McLachlan, A., Winter, P.E.D. & De Ruyek, A.M.C. (1998). Impact of trampling on sandy beach macrofauna. Journal of Coastal Conservation **4**: 87 – 90.
- Moss, D. & McPhee, D.P. (2006). The impacts of recreational four-wheel driving on the abundance of the ghost crab (*Ocypode cordimanus*) on a subtropical sandy beach in SE Queensland. Coastal Management **34**: 133 – 140.

- Nel, R., McLachlan, A. & Winter, D. (1999). The effect of sand particle size on the burrowing ability of the beach Mysid *Gastrosaccus psammodytes* Tattersall. Estuarine, Coastal and Shelf Science **48**: 599 – 604.
- Nel, R., McLachlan, A. & Winter, D.P.E. (2001). The effect of grain size on the burrowing of two *Donax* species. Journal of Experimental Marine Biology and Ecology **265**: 219 – 238.
- Nicholas, W.L. (2001). Seasonal variations in nematode assemblages on an Australian temperate ocean beach; the effect of heavy seas and unusually high-tides. Hydrobiologia **464**: 14 – 26.
- Nicholas, W.L. & Hodda, M. (1999). The free-living nematodes of a temperate, high-energy, sandy beach: faunal composition and variation of space and time. Hydrobiologia **394**: 22 - 127.
- Olabarria, C., Lastra, M. & Garrido, J. (2007). Succession of macrofauna on macroalgal wrack of an exposed sandy beach: Effects of patch size and site. Marine Environmental Research **63**: 19 – 40.
- Papageorgiou, N., Moreno, M., Marin, V., Baiardo, S., Avantidis, C., Fabiano, M. & Eleftheriou, A. (2007). Interrelationships of bacteria, meiofauna and macrofauna in a Mediterranean sedimentary beach (Maremma Park, NW Italy). Helgoland Marine Research **61**: 31 – 42.
- Platt, H.M. & Warwick, R.M. (1983). Freelifving marine nematodes. Part I: British enlopids. Synopsis of the British fauna no. 28. Cambridge University Press, Cambridge.
- Platt, H.M. & Warwick, R.M. (1988). Freelifving marine nematodes. Part II: British Chromadorida. Synopsis of the British fauna no. 38. Brill, Leiden.
- Polis, G.A. & Hurd, S.H. (1995). Extraordinarily high spider densities on islands: flow of energy from the marine to terrestrial food webs and the absence of predation. Proceedings of the National Academy of Sciences **92**: 4382 – 4386.
- Quinn, G.P. & Keough, M.J. (2002). Experimental design and data analysis for biologists. Cambridge University Press, Cambridge.
- Rodríguez, J.G., Lastra, M. & Lopez, J. (2003). Meiofauna distribution along a gradient of sandy beaches in northern Spain. Estuarine, Coastal and Shelf Science **58S**: 63 – 69.

- Rodríguez, J.G., López, J. & Jaramillo, E. (2001). Community structure of the intertidal meiofauna along a gradient of morphodynamic sandy beach types in southern Chile. Revista Chilena de Historia Natural **74**: 885 – 897.
- Salvat, B. (1964). Les conditions hydrodynamiques interstitielles des sédiments meubles intertidaux et la répartition verticale de la faune endogée. Académie des Sciences **259**: 1576 – 1579.
- Salvat, B. (1967). La macrofaune carinologique endogée des sédiments meubles intertidaux (tanaïdes, isopodes et amphipodes): éthologie, biologie et cycle biologique. Mémoires de la Muséum National d'Histoire Naturelle, Série A **45**: 1 – 275.
- Scapini, F. (2003). Beaches – what future? An integrated approach to the ecology of sandy beaches. Estuarine, Coastal and Shelf Science **58S**: 1 – 3.
- Schlacher, T.A. & Connolly, R.M. (2009). Land-ocean coupling of carbon and nitrogen fluxes on sandy beaches. Ecosystems **12**: 311 – 321.
- Schlacher, T.A. & Morrison, J.M. (2008). Beach disturbance caused by off-road vehicles (ORVs) in sandy shores: relationship with traffic volumes and a new method to quantify impacts using image-based acquisition and analysis. Marine Pollution Bulletin **56**: 1646 – 1649.
- Schlacher, T.A. & Thompson, L.M.C. (2007). Exposure of fauna to off-road vehicle (ORV) traffic on sandy beaches. Coastal Management **35**: 567 – 583.
- Schlacher, T.A. & Thompson, L.M.C. (2008). Physical impacts caused by off-road vehicles on sandy beaches: spatial quantification of car tracks on an Australian barrier island. Journal of Coastal Management **24**: 234 – 242.
- Schlacher, T.A., Dugan, J., Schoeman, D.S., Lastra, M., Jones, A., Scarpini, F., McLachlan, A. & Defeo, O. (2007a). Sandy beaches at the brink. Diversity and Distributions **13**: 556 – 560.
- Schlacher, T.A., Richardson, D. & McLean, I. (2008c). Impacts of off-road vehicles (ORVs) on macrobenthic assemblages on sandy beaches. Environmental Management **41**: 878 – 892.
- Schlacher, T.A., Schoeman, D.S., Dugan, J., Lastra, M., Jones, A., Scarpini, F. & McLachlan, A. (2008a). Sandy beach ecosystems: key features,

- sampling issues, management challenges and climate change impacts. Marine Ecology **29S**: 70 – 90.
- Schlacher, T.A., Schoeman, D.S., Lastra, M., Jones, A., Dugan, J., Scarpini, F., & McLachlan, A. (2006). Neglected ecosystems bear the brunt of change. Ethology, Ecology & Evolution **18**: 349 – 351.
- Schlacher, T.A., Thompson, L. & Price, S. (2007b). Vehicles versus conservation of invertebrates on sandy beaches: mortalities inflicted by off-road vehicles on ghost crabs. Marine Ecology **28**: 354 – 367.
- Schlacher, T.A., Thompson, L. & Walker, S.J. (2008b). Mortalities caused by off-road vehicles (ORVs) to a key member of sandy beach assemblages, the surf clam *Donax deltoides*. Hydrobiologia **610**: 345 – 350.
- Schooler, N., Dugan, J.E., Page, H.M. & Hubbard, D. (2009). Dispersal of talitrid amphipods, *Megalorchestia* spp. on sandy beaches: implications for ecological recovery following a disturbance. *In* 5<sup>th</sup> International Symposium on Sandy Beaches (Abstracts). University of Mohammed V Agdal, Scientific Institute, Rabat, Morocco, 19<sup>th</sup> – 23<sup>rd</sup> October, 2009.
- Schratzberger, M., Gee, J.M., Rees, H.L., Boyd, S.E. & Wall, C.M. (2000). The structure and taxonomic composition of sublittoral meiofauna assemblages as an indicator of the status of marine environments. Journal of the Marine Biological Association of the United Kingdom **80**: 969 – 980.
- Sheppard, N., Pitt, K.A. & Schlacher, T.A. (2009). Sub-lethal effects of off-road vehicles (ORVs) on surf clams on sandy beaches. Journal of Experimental Marine Biology and Ecology **380**: 113 – 118.
- Sherman, K.M. & Coull, B.C. (1980). The response of meiofauna to sediment disturbance. Journal of Experimental Marine Biology and Ecology **46**: 59 – 71.
- Shillabeer, N., Hart, B. & Riddle, A.M. (1992). The use of a mathematical model to compare particle size data derived by dry-sieving and laser analysis. Estuarine, Coastal and Shelf Science **35**: 105 – 111.
- Short, A.D. (1996). The role of wave height, period, slope, tide range and embaymentization in beach classifications: A review. Revista Chilena De Historia Natural **69**: 589 – 604.
- Short, A.D. (ed.) (1999). Handbook of beach and shore morphodynamics. Wiley and Sons, London.

- Short, A.D. (2006a). Beaches of the South Australian coast and Kangaroo Island: A guide to their nature, characteristics, surf and safety. Sydney University Press, Sydney.
- Short, A.D. (2006b). Australian beach systems - Nature and distribution. Journal of Coastal Research **22**: 11 – 27.
- Short, A.D. & Wright, L.D. (1983). Physical variability of sandy beaches. In: *Sandy beaches as ecosystems*. A. McLachlan & T. Erasmus. (eds.): pp. 133-144, The Hague: Dr W Junk.
- Short, A.D. & Wright, L.D. (1984). Morphodynamics of high energy beaches and surf zones: an Australian perspective. In: *Coastal geomorphology in Australia*. B.G. Thom (ed.). Academic Press, Sydney.
- Soares, A.G., Schlacher, T.A. & McLachlan, A. (1997). Carbon and nitrogen exchange between sandy beach clams (*Donax serra*) and kelp beds in the Benguela coastal upwelling region. Marine Biology **127**: 657 – 664.
- Somerfield, P.J. & Warwick, R.M. (1996). Meiofauna in marine pollution monitoring programmes. A laboratory manual. Ministry of Agriculture, Fisheries and Food, Directorate of Fisheries Research, Lowerstoft, UK.
- Somerfield, P.J., Warwick, R.M. & Moens, T. (2005). Meiofauna techniques. In: *Methods for the study of Marine Benthos*. 3rd edition. A. Eleftheriou and A. D. McIntyre (eds.). Blackwell Science, New Delhi, India. pp. 229 – 272.
- Steiner, A.J. & Leatherman, S.P. (1981). Recreational impacts on the distribution of ghost crabs *Ocypode quadrata* FAB. Biological Conservation **20**: 111 – 122.
- Stephens, E. (2004). Life's a beach: habitat preference and disturbance impacts on the hooded plover, a globally threatened shorebird. Unpublished Honours Thesis, University of Adelaide.
- Stephens, E. (2009). Assessment of Hooded Plover (*Thinornis rubricollis*) breeding success and threats on the Fluerieu Peninsula 2006/07. Report to the Hooded Plover Management Project.
- Ugolini, A., Ungherese, G., Somigli, S., Galanti, G., Baroni, D., Borghini, F., Cipriani, N., Nebbiai, M., Passaponti, M. & Focardi, S. (2008). The amphipod *Talitrus saltator* as a bioindicator of human trampling on sandy beaches. Marine Environmental Research **65**: 349 – 357.

- Underwood, A.J. (1997). Experiments in ecology: their logical design and interpretation using analysis of variance. Cambridge University Press, Cambridge.
- Van Dam, A.R. & Van Dam, M.H. (2008). Impact of off-road vehicle use on dune endemic Coleoptera. Annals of the Entomological Society of America **101**: 411 – 417.
- van der Merwe, D. & van der Merwe, D. (1991). Effects of off-road vehicles on the macrofauna of a sandy beach. South African Journal of Science **87**: 210 – 213.
- Veloso, V.G. & Cardoso, R.S. (2001). Effect of morphodynamics on the spatial and temporal variation of macrofauna on three sandy beaches, Rio de Janeiro State, Brazil. Journal of the Marine Biological Association of the United Kingdom **81**: 369 – 375.
- Ward, T.J. & Jacoby, C.A. (1992). A strategy for assessment and management of marine ecosystems: baseline and monitoring studies in Jervis Bay, a temperate Australian embayment. Marine Pollution Bulletin **25**: 163 – 171.
- Warwick, R.M. & Clarke, K.R. (1993). Increased variability as a symptom of stress in marine communities. *Journal of Experimental Marine Biology and Ecology* **172**: 215 – 226.
- Warwick, R.M., Platt, H.M. & Sommerfield, P.J. (1998). Freelifving marine nematodes. Part III: British Monhysterida. Synopsis of the British fauna no. 53. Field Studies Council, Shrewsbury.
- Webb, J.E. (1991). Hydrodynamics, organisms and pollution of coastal sands. Ocean & Shoreline Management **16**: 23 – 51.
- Webb, R.H. (1982). Off-road motorcycle effects on a desert soil. Environmental Conservation **9**: 197 – 208.
- Webb, R.H. (1983). Compaction of desert soils by off-road vehicles. In: *Environmental effects of off-road vehicles*. R.H. Webb and H.G. Wilshire (eds.). Springer-Verlag, New York.
- Weiser, W. (1953). Die Beziehung zwischen Mundhöhlengestalt, Ernährungsweise und Vorkommen bei freilbenden marinen Nematoden. Arkiv fur Zoologie **4**: 449 – 484.
- Wentworth, C.K. (1922). A scale of grade and class terms for clastic sediments. Journal of Geology **30**: 377 – 392.

- Wilshire, H.G., Nakata, J.K., Shipley, S. & Prestegard, K. (1978). Impacts of vehicles on natural terrain at seven sites in the San Francisco Bay area. Environmental Geology **2**: 295 – 319.
- Wolcott, T.G. & Wolcott, D.L. (1984). Impact of off-road vehicles on macroinvertebrates of a mid-Atlantic beach. Biological Conservation **29**: 217 – 240.
- Wright, L.D. & Short, A.D. (1983). Morphodynamics of beaches and surf zones in Australia. In: *CRC Handbook of Coastal Processes and Erosion*. P.D. Komar (ed.). pp. 35-64, CRC Press, Inc., Boca Raton.
- Wright, L.D. & Short, A.D. (1984). Morphodynamic variability of surf zones and beaches: a synthesis. Marine Geology **56**: 93 – 118.
- Yeates, G.W., Bongers, T., de Goede, R.G.M., Freckman, D.W. & Georgieva, S.S. (1993). Feeding habits in soil nematode Families and Genera – An outline for soil ecologists. Journal of Nematology **25**: 315 – 331.
- Zaremba, R., Godfrey, P.J. & Leatherman, S.P. (1979). The ecological effects of off-road vehicles on the beach/backshore (drift line) zone in Cape Cod National Seashore, Massachusetts, UM/NPSCRU report no. 29. Amherst, Mass.: The Environmental Institute, Univ. Massachusetts.